arXiv:0808.0158v1 [math.AG] 1 Aug 2008

APPROXIMATE ROOTS, TORIC RESOLUTIONS AND DEFORMATIONS OF A PLANE BRANCH

P.D. GONZÁLEZ PÉREZ

Abstract. We analyze the expansions in terms of the approximate roots of a Weierstrass polynomial $f \in \mathbf{C}\{x\}[y]$, defining a plane branch $(C, 0)$, in the light of the toric embedded resolution of the branch. This leads to the definition of a class of (non equisingular) deformations of a plane branch $(C, 0)$ supported on certain monomials in the approximate roots of f. As a consequence we find out a Kouchnirenko type formula for the Milnor number $(C, 0)$. Our results provide a geometrical approach to Abhyankar's straight line conditions and its consequences. As an application we give an equisingularity criterion for a family of plane curves to be equisingular to a plane branch and we express it algorithmically.

Introduction

The use of *approximate roots* in the study of plane algebraic curves, initiated by Abhyankar and Moh in [A-M], was essential in the proof of the famous *embedding line theorem* in [A-M2]. Let $(C, 0) \subset (\mathbf{C}^2, 0)$ be a germ of analytically irreducible plane curve, a *plane branch* in what follows. Certain *approximate roots* of the Weierstrass polynomial defining $(C, 0)$ are *semi-roots*, i.e., they define *curvettes* at certain exceptional divisors of the minimal embedded resolution. A'Campo and Oka describe the embedded resolution of a plane branch by a sequence of toric modifications using approximate roots in [A'C-Ok] and give topological proofs of some of the results of Abhyankar and Moh. See [Abh3], [PP], [G-P] or [As-B] for an introduction to the notion of approximate root and its applications.

We consider at the infinitely near points of the toric embedded resolution the canonical local coordinates defined by the strict transform of a suitable approximate root (or semi-root) and the exceptional divisor. In Section 2 we introduce an injective correspondence between monomials in these coordinates and monomials in the approximate roots (see Proposition 2.1). From this natural correspondence we derive two applications.

The first one is the definition of a class of (non equisingular) multi-parametric deformations $C_{\underline{t}}$ of the plane branch, which we call multi-semi-quasi-homogeneous (msqh). We explain its basic properties in Section 3. The terms appearing in this deformation are monomials in the approximate roots of f and the deformation may be seen naturally as a deformation of Teissier's embedding of the plane branch C in a higher dimensional affine space (see [T3]). If the deformation $C_{\underline{t}}$ is *generic* the Milnor number of $(C, 0)$ is related to the sum of the Milnor numbers of some curves defined from $C_{\underline{t}}$ at the infinitely near points of the toric resolution of $(C, 0)$ (see Proposition 36). As a consequence we derive a Kouchnirenko type formula for the Milnor number in terms of certain polyhedral complex with integral structure which we associate to $(C, 0)$. In a joint work with Risler we apply the construction of this class of deformations in the study of the topological types of smoothings of real plane branches with the maximal number of connected components (see [GP-R]).

2000 *Mathematics Subject Classification.* Primary 14J17; Secondary 32S10, 14M25.

Key words and phrases. approximate roots, deformations of a plane curve, equisingularity criterion.

Supported by *Programa Ramón y Cajal* and by MTM2007-6798-C02-02 grants of *Ministerio de Educación y Ciencia*, Spain.

The second application, given in Section 4, is based on the relations of the expansions in terms of approximate-roots, Teissier's embedding of the plane branch and Abhyankar's *straight line condition* for the *generalized Newton polygons* associated to a plane branch. These relations are better understood by passing through the toric embedded resolution of the branch (see Theorem 4.1 and Proposition 1.16). In particular, we prove that the generalized Newton polygons arise precisely from the Newton polygons of the strict transform of $(C, 0)$ at the infinitely near points of the toric embedded resolution of $(C, 0)$ (see Proposition 4.5). We have revisited Abhyankar's irreducibility criterion for power series in two variables (see [Abh4]). We give a proof of Abhyankar's criterion by using the toric geometry tools we have previously introduced. As an application we obtain an equisingularity criterion for a family of plane curves to be equisingular to a plane branch (see Theorem 4.3), which generalizes the one given by A'Campo and Oka in [A'C-Ok]. The criterion is expressed as algorithmic procedure (see Algorithm 4.7). Alternatively, we can use Algorithm 4.8 which is derived independently from the irreducibility criterion of García Barroso and Gwoździewicz in [GB-G], in terms of jacobian Newton polygons.

The paper is organized as follows: Section 1 introduce basic results and definitions. Section 3 only depends on Section 1 and 2. Section 4 do not depend on Section 3.

Notations. We denote by $[\alpha]$ the integer part of $\alpha \in \mathbf{Q}$. We denote by $\deg P$ (resp. by $\deg_y P$) the degree of a polynomial P (resp. with respect to the indeterminate y). The ring of formal (resp. convergent) power series in $\underline{x} = (x_1, \dots, x_m)$ is denoted by $\mathbf{C}[[\underline{x}]]$ (resp. by $\mathbf{C}\{\underline{x}\}$). If $0 \neq \alpha \in \mathbf{C}[[t]]$ is a series, the order $\mathrm{ord}_t \alpha$ of α is the integer n such that $\alpha = t^n \epsilon$, where $\epsilon \in \mathbf{C}[[t]]$ and $\epsilon(0) \neq 0$; the number $\epsilon(0)$ is called the initial coefficient of α. If $f = \sum c_{i,j} x^i y^j \in \mathbf{C}[[x, y]]$ is a series, the Newton polygon $\mathcal{N}(f)$ is the convex hull of the set $\bigcup_{c_{i,j} \neq 0} (i, j) + \mathbf{R}^2_{\geq 0}$. If $\Lambda \subset \mathbf{R}^2$ the symbolic restriction of f to Λ is $\sum_{(i,j) \in \Lambda \cap \mathbf{Z}^2_{\geq 0}} c_{i,j} x^i y^j$. If $(C_i, 0) \subset (\mathbf{C}^2, 0)$, $i = 1, 2$ are plane curve germs defined by $h_i(x, y) = 0$, for $h_i \in \mathbf{C}\{x, y\}$, we denote by $(C_1, C_2)_0$ or by $(h_1, h_2)_0$ the intersection multiplicity $\dim_{\mathbf{C}} \mathbf{C}\{x, y\}/(h_1, h_2)$.

1. Plane branches, semi-roots and toric resolution

We introduce first our notations and we review some well known invariants associated to a plane branch $(C, 0)$ embedded in $(\mathbf{C}^2, 0)$. For more details see for instance [Z2], [W], [PP], [T3], [Abh3], [C], and [T4].

1.1. Equisingularity invariants of a plane branch.

We consider an analytically irreducible plane curve germ, a plane branch for short, $(C, 0) \subset (\mathbf{C}^2, 0)$. Let (x, y) be a pair of local coordinates at $(\mathbf{C}^2, 0)$ such that the germ of smooth curve $(L, 0)$ defined by $x = 0$ has intersection multiplicity $(C, L)_0 = n < \infty$. By the Weierstrass Preparation Theorem $(C, 0)$ is defined by an equation $f(x, y) = 0$ where $f \in \mathbf{C}\{x\}[y]$ is a *Weierstrass polynomial* in y. By Newton Puiseux Theorem the branch $(C, 0)$ has a fractional power series parametrization of the form, $y = \zeta$, where:

$$\zeta = \sum_i a_i x^{i/n}, \quad a_i \in \mathbf{C}, \tag{1}$$

and $n = \deg_y f$. We set $e_0 := n$ and then inductively for $j \geq 1$, b_j the smallest integer $i > b_{j-1}$ not divisible by e_{j-1} with $a_i \neq 0$, $e_j := \gcd(e_{j-1}, b_j)$. We obtain in this way a finite sequence which stops when we reach an integer g such that $e_g = \gcd(e_{g-1}, b_g) = 1$. We obtain two sequences of integers $e_0, \dots, e_g = 1$ and $n_0 := 1, n_1 := e_0/e_1, \dots, n_g = e_{g-1}/e_g$, where $n_j > 0$ for $j = 1, \dots, g$ (see [Z2]).

Definition 1.1. (see [Z1], pag. 993) *The sequence* $b_1/n, \dots, b_g/n$ *(resp.* $n_1, \dots, n_g$*) is called sequence of characteristic exponents (resp. characteristic integers) of* $(C, 0)$ *with respect to* L.

This sequence does not depend on the possible choice of local coordinate y once x has been fixed. If L is not tangent to C, then b_0 is the multiplicity of $(C, 0)$ and the characteristic exponents b_j relative to L, also denoted by β_j in the literature, do not depend on the choice of non tangent curve L. This sequence is equivalent to the sequence of classical characteristic pairs, which classifies the equisingularity class of the branch (see [Z1] and [Abh1]).

1.1.1. *The semigroup of a branch.* The map of germs $(\mathbf{C}, 0) \to (C, 0)$, given by $t \mapsto (x(t) := t^n, y(t) := \zeta(t))$, (which defined by replacing $x^{1/n}$ by t in (1)) is the *normalization* of the branch $(C, 0)$. It corresponds algebraically to the inclusion

$$\mathcal{O}_C \to \bar{\mathcal{O}}_C = \mathbf{C}\{t\}, \quad h \mod (f) \mapsto h(x(t), y(t)). \tag{2}$$

of the local ring of the branch in its *integral closure in its field of fractions.*

If $h(x, y) \in \mathbf{C}\{x\}[y]$ defines a plane curve germ, we have that $(f, h)_0 = \mathrm{ord}_t(h(x(t), y(t))$.

The set $\Lambda := \{(f, h)_0 \mid h \in \mathbf{C}\{x, y\} - (f)\}$ is a sub-semigroup of $\mathbf{Z}_{\geq 0}$ of finite type associated to the plane branch C. The semigroup Λ is generated by:

$$\bar{b}_0 := e_0, \quad \bar{b}_1 := b_1, \quad \bar{b}_{j+1} := n_j \bar{b}_j + b_{j+1} - b_j \quad \text{for } j = 1, \dots, g-1. \tag{3}$$

We set $\bar{b}_{g+1} := \infty$. The generators (3) are called the *characteristic sequence of generators* of the semigroup Λ with respect to the line $x = 0$.

Lemma 1.2. *(see* [T3]*) Any* $\bar{b} \in \Lambda$ *has a unique expansion of the form:*

$$\bar{b} = \eta_0 \bar{b}_0 + \eta_1 \bar{b}_1 + \cdots + \eta_g \bar{b}_g, \tag{4}$$

where $0 \leq \eta_0$ *and* $0 \leq \eta_j < n_j$*, for* $j = 1, \dots, g$*. The image of* $\bar{b}_j$ *in the group* $\mathbf{Z}/(\sum_{i=0}^{j-1} \mathbf{Z}\bar{b}_i)$ *is of order* n_j*. We have that:*

$$n_j \bar{b}_j \in \mathbf{Z}_{\geq 0} \bar{b}_0 + \cdots + \mathbf{Z}_{\geq 0} \bar{b}_{j-1} \text{ and } n_j \bar{b}_j < \bar{b}_{j+1}, \text{ for } j = 1, \dots, g. \tag{5}$$

Remark 1.3. *Let us consider a semigroup* Θ *of* $\mathbf{Z}_{\geq 0}$ *with a minimal set of generators* $\bar{b}_0 < \cdots < \bar{b}_g$*. Set* $e_0 := \bar{b}_0$*,* $e_j := \gcd(\bar{b}_0, \dots, \bar{b}_j)$ *and* $n_j := e_{j-1}/e_j$ *for* $j = 1, \dots, g$*. Then, if conditions (5) hold the semigroup* Θ *is the semigroup associated with a plane branch (see* [T3]*).*

1.2. **Approximate roots, semi-roots and truncations.** For an introduction to these topics see the survey [PP] of Popescu-Pampu. See also [Abh4], [Z2], [A'C-Ok], and [G-P] and [LJ].

1.2.1. *Approximate roots.* Let A be a integral domain containing $\mathbf{Q}$. Let $H \in A[y]$ be a monic polynomial in y of degree > 0. Any polynomial $F \in A[y]$ has a unique H*-adic expansion* of the form:

$$F = a_s + a_{s-1} H + \cdots + a_1 H^{s-1} + a_0 H^s, \tag{6}$$

where $a_i \in A[y]$, $\deg a_i < \deg H$ and $s = [\deg F / \deg H]$ (for $[a]$ denoting integral part of a). This expansion is obtained by iterated Euclidean division by H (see [Z2]).

Denote by $\mathcal{B}_m \subset A[y]$ the set of monic polynomials of degree $m > 0$ in y. Let $F \in A[y]$ be a monic polynomial of degree N divisible by m. Set $N = mk$. The *Tschirnhausen operator* $\tau_F : \mathcal{B}_m \to \mathcal{B}_m$ is defined by $\tau_F(H) = H + \frac{a_1}{k}$ where a_1 is the coefficient of H^{k-1} in the H-adic expansion (6) of F. For instance, if $m = 1$, $H = y$ and $y' :== y + \frac{a_1}{N}$, then the coefficient of $(y')^{N-1}$ in the y'-expansion of F is zero. Setting $y' = \tau_F(y)$ defines a change of coordinates, which is classically called the *Tschirnhausen transformation.*

Definition 1.4. *Let A a domain containing $\mathbf{Q}$. Let $F \in A[y]$ a monic polynomial of degree N and suppose $N = mk$. An approximate root G of degree m of F is a monic polynomial in $A[y]$ such that* $\deg(F - G^k) < N - m$.

The approximate root G of degree m of F exists and is unique. It is determined algorithmically in terms of Euclidean division of polynomials by: $G = \tau_F \circ \overset{(m)}{\cdots} \circ \tau_F(H), \quad \forall\, H \in \mathcal{B}_m$.

Lemma 1.5. *Let A be an integral domain containing $\mathbf{Q}$ and $F \in A[[x]][y]$ a polynomial of the form: $F = (y^p + cx^q)^e + \cdots$ where $c \in A \setminus \{0\}$ and the exponents of the terms which are not written lie in the interior the Newton polygon of F. Let m be a positive integer dividing e. Then, the approximate root G of degree mp of the form: $G = (y^p + cx^q)^m + \cdots$, where the exponents of the terms which are not written lie in the interior of the Newton polygon of G.*

Proof. Let $v = (p, q)$ be the normal vector to the compact edge of the Newton polygon of F. For $H \in A[[x]][y]$ we denote by $H_{|v}$ the symbolic restriction of H to the compact face of its Newton polygon determined by v (see [G-T]). If we give weight p, q to x, y respectively, we find that $F_{|v}$ is quasi-homogeneous of weight pqe and $F = F_{|v} + \cdots$, where the dots means terms of weight $> epq$. It follows that $G_{|v} = y^{mp} + \cdots$ is quasi-homogeneous of weight mpq and $G = G_{|v} + \cdots$, where the dots means terms of weight $> mpq$. Since G is an approximate root we have that $\deg_y(F - G^{e/m}) < \deg_y F - \deg_y G$. Using the weights we deduce that: $\deg_y(F_{|v} - G_{|v}^{e/m}) < \deg_y F_{|v} - \deg_y G_{|v}$. This implies that $G_{|v}$ is an approximate root of $F_{|v}$ of degree mp. By the unicity of the approximate roots it follows that $G_{|v} = (y^p + cx^q)^m$. □

1.2.2. *Semi-roots.*

Definition 1.6. *Let $(C, 0) \subset (\mathbf{C}^2, 0)$ be a plane branch and $(L, 0) \subset (\mathbf{C}^2, 0)$ be a germ of smooth curve. A j-semi-root $(C_j, 0)$ of $(C, 0)$ with respect to L, is a germ of curve such that*

$$(C_j, C)_0 = \bar{b}_j \text{ and } (C_j, L)_0 = n_0 \cdots n_{j-1}, \quad if\, 0 \leq j \leq g, \tag{7}$$

where $\bar{b}_j$ and n_j are respectively the generators of the semigroup (3) and the characteristic integers, with respect to L. We set $C_{g+1} = C$.

The plane curve singularity parameterized by, $y = \zeta_j$, where $\zeta_j = \sum_{i<b_j} a_i x^{i/n}$ is the fractional power series obtained from (1) by truncation $\mod x^{b_j/n}$. It is a j-semi-root of $(C, 0)$ with respect to $x = 0$ (see [Z2]). Notice that ζ_j belongs to $\mathbf{C}\{x^{1/n_1 \cdots n_{j-1}}\}$.

Proposition 1.7. (see [PP], [Z2]) *The following assertions are equivalent:*

1. *The germ $(D, 0)$ is a j-semi-root of $(C, 0)$ with respect to the line L, given by $x = 0$.*
2. *The germ $(D, 0)$ is a plane branch with $(D, L) = n_1 \cdots n_{j-1}$ and in the local coordinates (x, y) it is parameterized by a fractional power series $y = \tau(x^{1/n_1 \cdots n_{j-1}})$ such that $\tau = \zeta \mod x^{b_j/n}$, where ζ is the parametrization (1) of $(C, 0)$.*

It follows that in the coordinates (x, y) a j-semi-root can be defined by a Weierstrass polynomial $f_j \in \mathbf{C}\{x\}[y]$ of degree $n_0 \cdots n_j$ in y.

The notion of semi-root is related with Lejeune-Jalabert's notion of curves of maximal contact with higher genus, which is defined also over fields of arbitrary characteristic [LJ]. The notion of semi-root is also related with that of approximate root by the following Theorem of Abhyankar and Moh (see [A-M], [G-P] and [PP]).

Proposition 1.8. *Let $f \in \mathbf{C}\{x\}[y]$ be a Weierstrass polynomial defining a plane branch $(C, 0)$. The approximate roots of f of degrees $n_0 \cdots n_{j-1}$ define semi-roots of $(C, 0)$ with respect to $x = 0$, for $j = 1, \ldots, g$.*

1.2.3. *Expansions.*

Proposition 1.9. (see [Abh2] and [PP]) *Let us consider a sequence of integers* $n_1, \dots, n_g > 1$. *If* $F_1, \dots, F_{g+1} \in \mathbf{C}\{x\}[y]$ *are polynomials of degrees* $1, n_1, n_1 n_2, \dots, n_1 \cdots n_g$, *respectively. then any polynomial* $F \in \mathbf{C}\{x\}[y]$ *has a unique expansion of the form:*

$$F = \sum_I \alpha_I(x) F_1^{i_1} \cdots F_g^{i_g} F_{g+1}^{i_{g+1}}, \text{ with } \alpha_I(x) \in \mathbf{C}\{x\}, \tag{8}$$

where the components of the index $I = (i_1, \dots, i_{g+1})$ *verify that* $0 \leq i_1 < n_1, \dots, 0 \leq i_g < n_g$, $0 \leq i_{g+1} \leq [\deg_y F / \deg_y F_{g+1}]$. *Moreover, the degrees in* y *of the terms* $F_1^{i_1} \cdots F_{g+1}^{i_{g+1}}$ *are all distinct.*

Proof. Consider the F_{g+1}-adic expansion, of the form (6), of the polynomial F. Iterate the procedure by taking recursively F_j-adic expansions of the coefficients obtained, for $j = 1, \dots, g$ in decreasing order. The assertion of the degrees in y is consequence of the following elementary property of the sequence of integers $(n_1, \dots, n_g)$ (see [PP], proof of corollary 1.5.4). ☐

Remark 1.10. *With the above notations set*

$$\mathcal{A}_{g+1} := \{ I = (i_1, \dots, i_{g+1}) \mid 0 \leq i_1, < n_1, \dots, 0 \leq i_g, < n_g, 0 \leq i_{g+1} \}.$$

Then, the map $\mathcal{A}_{g+1} \to \mathbf{Z}$, *given by* $I \mapsto q_I := i_1 + n_1 i_2 + \cdots + n_1 \cdots n_g i_{g+1}$, *is injective.*

By expanding the coefficients of (8) as a series in x we obtain an expansion of the form:

$$F = \sum_I a_I x^{i_0} F_1^{i_1} \cdots F_g^{i_g} F_{g+1}^{i_{g+1}}, \text{ with } a_I \in \mathbf{C}, \tag{9}$$

Definition 1.11. *We call the expansion (8) (resp. (9)) the* $(F_1, \dots, F_{g-1}, F_{g+1})$*-adic expansion (resp.* $(x, F_1, \dots, F_{g-1}, F_{g+1})$*-adic expansion) of* F.

Let us fix a sequence of semi-roots $(C_j, 0)$ of the plane branch $(C, 0)$, where each curve C_j is defined by a Weierstrass polynomial $f_j(x, y)$ of degree $n_0 \cdots n_{j-1}$, which we call also semi-root, for $j = 1, \dots, g+1$. We have that $f := f_{g+1}$ by definition.

Remark 1.12. *Notice that* $(C_k, 0)$ *for* $k = 1, \dots, j$, *is a sequence of semi-roots of* $(C_j, 0)$ *by Proposition 1.7.*

Since any element in the analytic algebra $\mathcal{O}_C$ of $(C, 0)$ is represented by the class, modulo (f), of a polynomial $F \in \mathbf{C}\{x\}[y]$ of degree $< n$ we can apply Proposition 1.9 in terms of the sequence of semi-roots and Lemma 1.2 to obtain the following formula for the intersection multiplicity:

$$(f, F)_0 = \min\{ (f, x^{i_0} f_1^{i_1} \cdots f_g^{i_g})_0 / c_I \neq 0 \}.$$

Notice that the number $(f, x^{i_0} f_1^{i_1} \cdots f_g^{i_g})_0 = \sum_{j=0}^{g} i_j \bar{b}_j \in \Lambda$ is expanded in the form (4), hence by Lemma 1.2 if $I \neq I'$, then the corresponding elements in the semigroup Λ are different.

Proposition 1.13. ([T3], [GP] Lemma 35) *If* $f_1, \dots, f_g$ *define a system of semi-roots we have the expansions of the form indicated in Proposition 1.13:*

$$\xi_{j+1} f_{j+1} = f_j^{n_j} - \theta_{j+1} x^{\eta_0^{(j)}} f_1^{\eta_1^{(j)}} \cdots f_{j-1}^{\eta_{j-1}^{(j)}} + \sum_{I_j = (i_0, \dots, i_j)} \alpha_{I_j} x^{i_0} f_1^{i_1} \cdots f_j^{i_j}, \tag{10}$$

such that $\theta_{j+1} \neq 0$ *and the integer*

$$\bar{b}(I_j) := i_0 \bar{b}_0 + \cdots + i_j \bar{b}_j \tag{11}$$

is expanded in the form (4), whenever $\alpha_I^{(j)} \neq 0$.

Moreover, if $\alpha_I^{(j)} \neq 0$ *we have that the inequality* $\bar{b}(I_j) \geq n_j \bar{b}_j$ *and equality holds only for the index* $I_0^j := (\eta_0^{(j)}, \dots, \eta_{j-1}^{(j)}, 0)$.

1.3. Semi-roots and the monomial curve.

In the following definition we fix an additional property for the sequence of semi-roots which simplify some of the equations and procedures defined below.

Definition 1.14. *The sequence* $(f_1, \dots, f_g)$ *is* normalized *if the image by (2) of* $f_j \mod (f)$ *in* $\mathbf{C}\{t\}$ *is of the form:*

$$f_j(x(t), y(t)) = t^{\bar{b}_j} \epsilon_j \in \mathbf{C}\{t\} \quad \text{with } \epsilon_j(0) = 1, \quad \text{for } j = 1, \dots, g. \tag{12}$$

Obviously, if $f_1, \dots, f_g$ defines a sequence of semi-roots we can normalize them by replacing f_j by $\eta_j f_j$ for some suitable constant $\eta_j \neq 0$. The statement of Proposition 1.9 remains true after these changes, by changing eventually the coefficients of the expansions.

Remark 1.15. *If the sequence of semi-roots is normalized then Proposition 1.13 holds with* $\theta_{j+1} = 1$ *in the expansion (10).*

Following Teissier and Goldin, we re-embed the pair $(C, 0) \subset (\mathbf{C}^2, 0)$ in the affine space $(\mathbf{C}^{g+1}, 0)$ using a sequence of normalized semi-roots. We re-embed first $(\mathbf{C}^2, 0)$ as a germ of smooth surface by

$$\mathbf{C}\{v_0, v_1, \dots, v_g\} \longrightarrow \mathbf{C}\{x, y\} \quad v_0 = x,\ v_1 = y,\ v_2 = f_1,\ \dots\ v_g = f_{g-1}. \tag{13}$$

This embedding induces also a re-embedding of $(C, 0)$ in $(\mathbf{C}^{g+1}, 0)$. Notice that the valuation of the coordinates induced by the parametrization of $(C, 0)$ induced natural weights of the coordinates $v_0, \dots, v_g$ given by the corresponding valuations $\bar{b}_0, \bar{b}_1, \cdots, \bar{b}_g$. In general the weight of a monomial $v^I = v_0^{i_0} v_1^{i_i} \cdots v_g^{i_g}$ is the integer $\bar{b}(I) = \sum i_j \bar{b}_j \in \Lambda$.

Proposition 1.16. ([T3], [G-T]) *Relations (10) correspond to a system of generators of the ideal of* $(C, 0) \subset (\mathbf{C}^{g+1}, 0)$. *The first* $g - 1$ *relations correspond to the embedding of* $(\mathbf{C}^2, 0) \subset (\mathbf{C}^{g+1}, 0)$.

Teissier proved Proposition 1.16 in [T3] by exhibiting a one-parameter deformation which has the plane branch $(C, 0)$ as generic fiber, while the special fiber is the *monomial curve* $C^\Lambda \subset \mathbf{C}^{g+1}$, defined by: $v_j^{n_j} - v_0^{\eta_0^{(j)}} v_1^{\eta_1^{(j)}} \cdots v_{j-1}^{\eta_{j-1}^{(j)}} = 0$, for $j = 1, \dots, g$. This deformation corresponds algebraically to the deformation of $\mathcal{O}_C$ into the associated graded ring $\mathrm{gr}_t \mathcal{O}_C$. Notice that the branch $(C, 0)$ and the monomial curve $(C^\Lambda, 0)$ have the same semigroup. The form of the equations is then described from the *miniversal deformation with constant semigroup* of the monomial curve (see [T3], [G-T] and [CN]). See also [GP] for a generalization of part of these results to the case of quasi-ordinary hypersurface singularities.

1.4. Embedded resolution of a plane branch by a sequence of toric morphisms.

An embedded resolution of the germ $(C, 0) \subset (\mathbf{C}^2, 0)$ is a modification $\pi : Z \to \mathbf{C}^2$ such that Z is a smooth surface, π is an isomorphism outside $0 \in \mathbf{C}^2$ and the pull back of $(C, 0)$ by π is a normal crossing divisor. An embedded resolution of $(C, 0)$ is classically obtained by blowing up 0 and the points of non transversality of the pull back of $(C, 0)$ in successive blow ups (see [Z1]). A branch $(D, 0) \subset (\mathbf{C}^2, 0)$ is called a *curvette* for the modification $\pi : Z \to \mathbf{C}^2$ if the strict transform of $(D, 0)$ by π is a smooth curve transversal to the exceptional divisor $E = \pi^{-1}(0)$ at a smooth point of E.

We recall the construction of an embedded resolution of a plane branch by a sequence of local toric modifications in terms of the characteristic exponents with respect to the line

$x = 0$. See [A'C-Ok], [Ok2], [GP] §3.4. See also [L-Ok], [Ok1], and [G-T] for more on toric geometry and singularities.

We define inductively a sequence of proper birational maps $\Pi_j : Z_{j+1} \to Z_j$, for $j = 1, \dots, g$. We fix a sequence $C_1, \cdots, C_g$ of semi-roots of C. The semi-roots C_j appear as curvettes in the embedded resolution and play an auxiliary role in the definition of the maps Π_j.

By convenience we denote $(\mathbf{C}^2, 0)$ by (Z_1, o_1). The semi-root C_1 together with the line $x = 0$ define a pair of local coordinates at the origin. Namely, we take local coordinates $(x_1 := x, y_1 := f_1)$ at o_1, where f_1 is a Weierstrass polynomial of degree 1 in y defining a semi-root C_1 of $(C, 0)$. By convenience we denote in the same way by $f_j(x_1, y_1)$ the polynomials in y_1 defining the semi-root C_j, for $j = 2, \dots, g+1$. By Proposition 1.7 the fractional power series expansion $y(x)$ of C, and also of C_j for $2 \leq j \leq g$ begins with the term $x^{b_1/n} = x^{m_1/n_1}$ where $m_1 := b_1/n_2 \cdots n_g$. It follows that

$$f(x_1, y_1) = (y_1^{n_1} - \theta_2 x_1^{m_1})^{e_1} + \cdots, \tag{14}$$

such that $\theta_2 \in \mathbf{C}^*$, gcd $(n_1, m_1) = 1$ and the integer $e_0 := e_1 n_1$ is the intersection multiplicity of C with the line, $x = 0$. The terms which are not written in (14) have exponents (i, j) such that $in_1 + jm_1 > n_1 m_1 e_1$, i.e., they lie above the compact edge

$$\Gamma_1 := [(0, n_1 e_1), (m_1 e_1, 0)]$$

of the Newton polygon of f. Notice that n_1, e_1 are the integers of the previous section.

Remark 1.17. *By Remark 1.12 we have that $f_j(x_1, y_1)$ is of the form indicated in the right hand side of Formula (14), by replacing e_1 by $e_1/n_j \cdots n_g$, for $j = 2, \dots, g$.*

The vector $\vec{p}_1 = (n_1, m_1)$ is orthogonal to Γ_1 and defines a subdivision of the positive quadrant $\mathbf{R}^2_{\geq 0}$, which is obtained by adding the ray $\vec{p}_1 \mathbf{R}_{\geq 0}$. The quadrant $\mathbf{R}^2_{\geq 0}$ is subdivided in two cones, $\tau_i := \vec{e}_i \mathbf{R}_{\geq 0} + \vec{p}_1 \mathbf{R}_{\geq 0}$, for $i = 1, 2$ and $\vec{e}_1, \vec{e}_2$ the canonical basis of $\mathbf{Z}^2$. We define the *minimal regular subdivision* Σ_1 of $\mathbf{R}^2_{\geq 0}$ which contains the ray $\vec{p}_1 \mathbf{R}_{\geq 0}$ by adding the rays defined by those integral vectors in $\mathbf{R}^2_{>0}$, which belong to the boundary of the convex hull of the sets $(\tau_i \cap \mathbf{Z}^2) \backslash \{0\}$, for $i = 1, 2$. We denote by σ_1 the unique cone of Σ_1 of the form, $\sigma_1 := \vec{p}_1 \mathbf{R}_{\geq 0} + \vec{q}_1 \mathbf{R}_{\geq 0}$ where $\vec{q}_1 = (c_1, d_1)$ satisfies that:

$$c_1 m_1 - d_1 n_1 = 1. \tag{15}$$

The subdivision Σ_1 defines a proper birational map $\Pi_1 : Z_2 \to Z_1$, which is obtained by gluing monomial maps $\mathbf{C}^2(\sigma) \to \mathbf{C}^2$, where σ runs through the set of two dimensional cones in Σ_1. For instance, the map $\pi_1 : \mathbf{C}^2(\sigma_1) \to \mathbf{C}^2$ is defined by:

$$\begin{array}{rcl} x_1 & = & u_2^{c_1} x_2^{n_1}, \\ y_1 & = & u_2^{d_1} x_2^{m_1}, \end{array} \tag{16}$$

where u_2, x_2 are canonical coordinates for the affine space $\mathbf{C}^2(\sigma_1)$. It should be noticed that the map Π_1 is a composition of point *blow ups*, as many as rays added in Σ_1 to subdivide $\mathbf{R}^2_{\geq 0}$. Each ray $\vec{a}\mathbf{R}_{\geq 0} \in \Sigma_1$ corresponds bijectively to a projective line $\mathbf{C}P^1$, embedded as an irreducible component of the exceptional divisor $\Pi_1^{-1}(0)$. We denote by $E_2 \subset Z_2$ the exceptional divisor defined by $x_2 = 0$ in the chart $\mathbf{C}^2(\sigma_1)$.

By convenience, we denote f by $f^{(1)}$ and $(C, 0)$ by $(C^{(1)}, o_1)$. We have that $\Pi_1^*(C^{(1)})$ defines a *Cartier divisor* on Z_2. For instance, on $\mathbf{C}^2(\sigma_1)$ it is defined by $f^{(1)} \circ \pi_1 = 0$, where $f^{(1)} \circ \pi_1$ decomposes as

$$f^{(1)} \circ \pi_1 = \mathrm{Exc}(f^{(1)}, \pi_1) \bar{f}^{(2)}(x_2, u_2), \text{ with } \bar{f}^{(2)}(0,0) \neq 0. \tag{17}$$

The *exceptional divisor* of $\Pi_1^*(C^{(1)})$ on this chart is defined by $\mathrm{Exc}(f^{(1)}, \pi_1) := y_1^{e_0} \circ \pi_1$.

The polynomial $\bar{f}^{(2)}(x_2, u_2)$ defines the *strict transform* $C^{(2)}$ of $C^{(1)}$, i.e., the closure of the pre-image by π_1^{-1} of the punctured curve $C^{(1)} \setminus \{o_1\}$ on the chart $\mathbf{C}^2(\sigma_1)$. We call $\mathrm{Exc}(F^{(1)}, \pi_1)$ the *exceptional divisor function* of $f^{(1)}$ on the chart defined by π_1. We analyze the intersection of the strict transform with the exceptional divisor on the chart $\mathbf{C}^2(\sigma_1)$: using (14) and (15) we find that $\bar{f}^{(2)}(x_2, 0) = 1$ and $\bar{f}^{(2)}(0, u_2) = (1 - \theta_2 u_2^{c_1 m_1 - d_1 n_1})^{e_1} = (1 - \theta_2 u_2)^{e_1}$. By a similar argument on the other charts it follows that E_2 is the only exceptional divisor of Π_1 which intersects the strict transform $C^{(2)}$, precisely at the point o_2 of the chart $\mathbf{C}^2(\sigma_1)$ with coordinates $x_2 = 0$ and $u_2 = \theta_2^{-1}$, with intersection multiplicity equal to e_1.

By Remark 1.17 we can apply this analysis for the semi-root C_2 and we deduce that the strict transform $C_2^{(2)}$ is a curvette for Π_1, i.e., more precisely, $C_2^{(2)}$ and the exceptional divisor E_2 have a normal crossing at the point o_2. The pair of germs $(E_2, C_2^{(2)})$ is defined by the pair of local coordinates (x_2, y_2), at the point $o_2 \in Z_2$. By Proposition 1.13 we have that:

$$y_2 = 1 - \theta_2 u_2 + x_2 R_2(x_2, u_2) \text{ for some } R_2 \in \mathbf{C}\{x_2, u_2\}. \tag{18}$$

The strict transform $C^{(2)}$ is defined in this coordinates by a function of the form:

$$f^{(2)}(x_2, y_2) = (y_2^{n_2} - \theta_3 x_2^{m_2})^{e_2} + \cdots, \tag{19}$$

where $\theta_3 \neq 0$, $\gcd(n_2, m_2) = 1$, $e_1 = e_2 n_2$ and the terms which are not written have exponents (i, j) such that $i n_2 + j m_2 > n_2 m_2 e_1$, i.e., they lie above the compact edge Γ_2 of the Newton polygon of $f^{(2)}(x_2, y_2)$. The result of substituting in $f^{(2)}(x_2, y_2)$, the term y_2 by using (18), is equal to $\bar{f}^{(2)}(x_2, u_2)$.

Proposition 1.18. (see [Z1], [A'C-Ok] and [GP] §3.4) *The characteristic exponents, characteristic integers of the branch $(C^{(2)}, o_2)$ with respect to the line E_2 are respectively:*

$$(e_1, \bar{b}_2 - n_1 \bar{b}_1, \ldots, \bar{b}_g - n_{g-1} \bar{b}_{g-1}), \quad (n_2, \ldots, n_g).$$

The strict transforms $C_2^{(1)}, \ldots, C_g^{(1)}$ of the sequence of semi-roots $C_2, \ldots, C_g$ of the plane branch $(C, 0)$ by π_1 define the sequence of semi-roots of the strict transform $C^{(1)}$ of $(C, 0)$.

Using Proposition 1.18 we can iterate this procedure defining for $j > 2$ a sequence of toric maps $\Pi_{j-1} : Z_j \to Z_{j-1}$, which are described by replacing the index 1 by $j - 1$ and the index 2 by j in the description above. In particular, when we refer to a Formula, like (15) at level j, we mean after making this replacement. Since by construction we have that $e_j | e_{j-1} | \cdots | e_1 | e_0$ (for $|$ denoting divides), at some step we reach a first integer g such that $e_g = 1$ and then the process stops. The composition of blow ups $\Pi_g \circ \cdots \circ \Pi_1$ is an *embedded resolution* of the germ $(C, 0)$.

Proposition 1.19. *We have the following statements for $j = 2, \ldots, g$: A j-semi-root C_j of $(C, 0)$ is a curvette by the component E_j of the exceptional divisor of* $\Pi_{j-1} \circ \cdots \circ \Pi_1$. *The strict transform $C_j^{(j)}$ of C_j is a smooth germ at $o_j \in C^{(j)}$ defining a* 1*-semi-root of $(C^{(j)}, o_j)$ with respect to the line E_j.*

Remark 1.20. *If the set of semi-roots is normalized then in the construction above $\theta_j = 1$, for $j = 2, \ldots, g$ and by (18) at level $\leq j$, the restriction of the function u_i to $x_j = 0$ is given by*

$$u_i = \begin{cases} 1 & \text{if} \quad i < j \\ 1 - y_j & \text{if} \quad i = j \end{cases} \tag{20}$$

We denote by $\mathrm{Exc}(f, \pi_1 \circ \cdots \circ \pi_j)$ the *exceptional function* defining the exceptional divisor of $(\Pi_1 \circ \cdots \circ \Pi_j)^*(C)$ on the chart $\mathbf{C}^2(\sigma_j)$. We have that

$$\mathrm{Exc}(f, \pi_1 \circ \cdots \circ \pi_j) = (y_1^{e_0} \circ \pi_1 \circ \cdots \circ \pi_j) \cdots (y_j^{e_j} \circ \pi_j). \tag{21}$$

Notation 1.21. *If* $p, q \in \mathbf{C}\{x, y\} \setminus (f)$ *define the meromorphic function* $h := p/q$ *we abuse of notation by denoting by* $\mathrm{ord}_t(h)$ *the integer* $\mathrm{ord}_t p(x(t), y(t)) - ord_t(q(x(t), y(t))$.

Lemma 1.22 provides the sequence of generators of the semigroup of the branch $(C^{(j)}, o_j)$ with respect to the line E_j.

Lemma 1.22. *We have that* $\mathrm{ord}_t(u_j) = 0$, $\mathrm{ord}(x_j) = e_{j-1}$ *and* $\mathrm{ord}_t f^{(j)}_{j+i} = \bar{b}_{j+i} - n_j \cdots n_{j+i-1}\bar{b}_j$, *for* $2 \leq j < g+1$ *and* $0 \leq i \leq g-j$. *Moreover, the following formula holds:*

$$\mathrm{ord}_t(\mathrm{Exc}(f, \pi_j \circ \cdots \circ \pi_1)) = e_{j-1}\bar{b}_j, \quad \text{for } 1 \leq j \leq g,$$

Proof. For $j = 2$, we deduce from (16) that $u_2 = x_1^{m_1} y_1^{-n_1}$. It follows from the definition that $\mathrm{ord}_t(u_1) = m_1 \mathrm{ord}_t(x_1) - n_1 \mathrm{ord}_t(y_1) = 0$. We have also that $x_1 = u_2^{c_1} x_2^{n_1}$ hence $\mathrm{ord}(x_2) = e_0/n_1 = e_1$. By (21) and (14) applied to f_{2+i} and π_1 we have that, $\mathrm{ord}_t(f_{2+i}) = \mathrm{ord}_t(y_1^{n_1 \cdots n_{2+i-1}}) + \mathrm{ord}_t(f^{(2)}_{2+i})$, for $0 \leq i \leq g-j$. Since $f^{(2)}_2 = y_2$ it follows that $\mathrm{ord}_t(y_2) = \bar{b}_2 - n_1\bar{b}_1$ and more generally $\mathrm{ord}_t(f^{(2)}_{2+i}) = \bar{b}_{2+i} - n_1 \cdots n_{i+1}\bar{b}_1$. The equality $\mathrm{ord}_t(\mathrm{Exc}(f, \pi_1) = e_0\bar{b}_1$, follows from (21). We conclude the proof by an easy induction on j using Proposition 1.18 and relations (3). □

2. Monomials determined from the embedded resolution

The following key Proposition shows that given $(r, s) \in \mathbf{Z}_{\geq 0}$ with $s < e_{j-1}$ there is a unique way to determine a suitable monomial $\mathcal{M}_j(r, s)$ in x_1 and the semi-roots $y_1, f_2, \ldots, f_j$ such that the composite $\mathcal{M}_j(r, s) \circ \pi_1 \circ \cdots \circ \pi_{j-1}$ is equal to the product of the exceptional divisor function $\mathrm{Exc}(f, \pi_1 \circ \cdots \circ \pi_{j-1})$ by the monomial $x_j^r y_j^s$ times a unit in the ring $\mathbf{C}\{x_j, y_j\}$.

To avoid cumbersome notations we denote simply by u_i the term $u_i \circ \pi_i \circ \cdots \circ \pi_{j-1}$, whenever $j > i$ is clear form the context.

Proposition 2.1. *If* $(r, s) \in \mathbf{Z}_{\geq 0}$ *with* $s < e_{j-1}$ *there exist unique integers* $k_2, \ldots, k_j, i_0, \ldots, i_j$ *such that*

$$0 < k_2, \ldots, k_j, \quad 0 < i_0, \quad 0 \leq i_1 < n_1, \quad \ldots, \quad 0 \leq i_{j-1} < n_{j-1}, \quad i_j = s, \tag{22}$$

verifying that

$$\mathrm{Exc}(f, \pi_1 \circ \cdots \circ \pi_{j-1})\, u_2^{k_2} \cdots u_j^{k_j}\, x_j^r\, y_j^s = (x_1^{i_0}\, y_1^{i_1}\, f_2^{i_2} \cdots f_j^{i_j}) \circ \pi_1 \circ \cdots \circ \pi_{j-1}. \tag{23}$$

The integers in Proposition 2.1 can be determined algorithmically by using the proofs of the following Lemmas.

Lemma 2.2. *If* $r \geq 0, l > 0$ *are integers there exist unique integers* k, i_0, i_1 *such that*

$$(y_1^{ln_1} \circ \pi_1) u_2^k x_2^r = x_1^{i_0} y_1^{i_1} \circ \pi_1 \text{ with } 0 < i_0, k \text{ and } 0 \leq i_1 < n_1.$$

More precisely, we have that:

$$k = l + [c_1 r/n_1], \quad i_0 = km_1 - rd_1, \text{ and } \quad i_1 = c_1 r - n_1[c_1 r/n_1] \tag{24}$$

In particular, $i_1 = 0$ *if and only if* $r = pn_1$ *for some integer* p.

Proof. By (16) we deduce that $u_2 = x_1^{m_1} y_1^{n_1}$ and $x_2 = x_1^{-d_1} y_1^{c_1}$. We have that the term $(y_1^{ln_1} \circ \pi_1) u_2^k x_2^r = (x_1^{km_1 - rd_1} y_1^{rc_1 + (l-k)n_1}) \circ \pi_1$ is the transform of a holomorphic monomial by π_1 if and only if: $0 \leq i_0' := km_1 - rd_1$ and $0 \leq i_1' := rc_1 + (l-k)n_1$, or equivalently, $\frac{d_1}{m_1} r \leq k \leq \frac{c_1}{n_1} r + l$. By (15) we have that $m_1 c_1 - d_1 n_1 = 1$. This implies that $\frac{d_1}{m_1} < \frac{c_1}{n_1}$, thus the interval of the real line $[\frac{d_1}{m_1} r, \frac{c_1}{n_1} r + l]$ is of length greater than $l \geq 1$. Any integer k lying on this interval is convenient to define a holomorphic monomial. The condition $i_1' < n_1$, is equivalent to $\frac{c_1}{n_1} r + l - k < 1$, and it is verified if and only if $k = [\frac{c_1}{n_1} r + l] = l + [\frac{c_1}{n_1} r] > 0$.

We denote the integers i'_0 and i'_1 corresponding to this choice of k by i_0 and i_1 respectively. We have then that:

$$i_0 = (\frac{c_1}{n_1} r + l) m_1 - r d_1 > (\frac{c_1}{n_1} r + l - 1) m_1 - r d_1 = r m_1 (\frac{c_1}{n_1} - \frac{d_1}{m_1}) + (l-1) m_1 \overset{(15)}{\geq} (l-1) m_1 \geq 0.$$

For the last assertion, we have that $i_1 = c_1 r - n_1 [c_1 r / n_1] = 0$ if and only if n_1 divides r, since $\gcd(c_1, n_1) = 1$ by (16). $\square$

Lemma 2.3. *If $(r,s) \in \mathbf{Z}_{\geq 0}$ with $s < e_1$ there exist unique integers k, i_0, i_1 with $0 < k, i_0$ and $0 \leq i_1 < n_1$ such that:* $\mathrm{Exc}(f, \pi_1) u_2^k {x_2}^r {y_2}^s = ({x_1}^{i_0} y^{i_1} f_2^s) \circ \pi_1$. *More precisely, we have that*

$$k = e_1 - s + [c_1 r / n_1], \quad i_0 = k m_1 - r d_1, \quad \text{and} \quad i_1 = c_1 r - n_1 [c_1 r / n_1]. \tag{25}$$

In particular, $i_1 = 0$ if and only if $r = p n_1$ for some integer p.

Proof. We use that $\mathrm{Exc}(f, \pi_1) = y_1^n \circ \pi_1$ by (21) and that $f_2^s \circ \pi_1 = (y_1^{s n_1} \circ \pi_1) {y_2}^s$. Hence we deduce that $\mathrm{Exc}(f, \pi_1) {y_2}^s = (y_1^{n - s n_1} f_2^s) \circ \pi_1$. Since $s < e_1$ we have that $n - s n_1 = n_1 (e_1 - s)$. Then we apply Lemma 2.2 for $r \geq 0$ and $l = e_1 - s > 0$. $\square$

Proof of Proposition 2.1. We prove the result by induction on the number g of characteristic exponents, with respect to the line $x_1 = 0$. The case $g = 1$ is proved in Lemma 2.3. By Proposition 1.18 the strict transform $C^{(2)}$ of the branch $C^{(1)}$ by π_1 has $g-1$ characteristic exponents with respect to the line $x_2 = 0$. By induction using (21), we have that if $(r,s) \in \mathbf{Z}^2_{\geq 0}$ and if $s < e_{j-1}$ there exist unique integers $k_3, \dots, k_j, i'_0, i_2, \dots i_j$ with $0 < i'_0$, $0 \leq i_2 < n_2, \dots,$ $0 \leq i_{j-1} < n_j$, $i_j = s$, and such that

$$\frac{\mathrm{Exc}(f, \pi_1 \circ \cdots \circ \pi_{j-1})}{(y^{e_0} \circ \pi_1 \cdots \circ \pi_{j-1})} u_3^{k_3} \cdots u_j^{k_j} x_j^r y_j^s = (x_2^{i'_0} y_2^{i_2} (f_3^{(1)})^{i_3} \cdots (f_j^{(1)})^{i_j}) \circ \pi_2 \cdots \circ \pi_{j-1}. \tag{26}$$

We show then that there exist unique integers $0 < k_2, i_0$ and $0 \leq i_1 < n_1$ such that

$$\mathrm{Exc}(f, \pi_1)\, u_2^{k_2} x_2^{i'_0} y_2^{i_2} (f_3^{(2)})^{i_3} \cdots (f_j^{(2)})^{i_j} = (x_1^{i_0}\, y_1^{i_1}\, f_2^{i_2} \cdots f_j^{i_j}) \circ \pi_1, \tag{27}$$

By (21), we have that: $\mathrm{Exc}(f, \pi_1)\, y_2^{i_2} (f_2^{(2)})^{i_2} \cdots (f_j^{(2)})^{i_j} = (y_1^q f_2^{i_2} \cdots f_j^{i_j}) \circ \pi_1$ where the integer q,

$$q = n_1 (e_1 - i_2 - n_2 i_3 - n_2 \cdots n_{j-1} i_j) = n_1 \left(n_2 (\cdots (n_{j-1} (e_{j-1} - i_j) - i_{j-1}) \cdots) - i_1\right) \tag{28}$$

is a positive multiple of n_1 by (22). Then we apply Lemma 2.2.$\square$

Definition 2.4. *If $0 \leq r$ and if $0 \leq s < e_{j-1}$ we define a monomial in $x, f_1, \dots, f_j$ by:*

$$\mathcal{M}_j(r,s) := x^{i_0} f_1^{i_1} \cdots f_j^{i_j} \tag{29}$$

by relation (23) in Proposition 2.1.

If $0 \leq r$ and if $0 \leq s < n_j$ we define similarly a monomial in $x, f_1, \dots, f_j$

$$\mathcal{M}_j^{(j+1)}(r,s) \tag{30}$$

by the using (29) when f is replaced by the semi-root f_{j+1}, for $j = 2, \dots, g-1$.

Remark 2.5. *Given $n_1 \dots n_j$ and the pairs $(n_1, m_1), \dots, (n_{j-1}, m_{j-1})$, the exponents appearing in the expansion of $\mathcal{M}_j^{(j+1)}(r,s)$ in terms of $x, f_1, \dots, f_{j-1}$ are determined algorithmically by (r,s) and conversely.*

Lemma 2.6. *If $0 \leq r$ and $s < e_{j-1}$, we have that:*

$$(\mathcal{M}_j(r,s), f)_0 = e_{j-2} \bar{b}_{j-1} + r e_{j-1} + s\, (\bar{b}_{j+1} - n_j \bar{b}_j), \quad \text{for } j = 2, \dots, g+1.$$

Proof. By Lemma 2.1 we have that:

$$\mathcal{M}_j(r,s)\circ\pi_1\circ\cdots\circ\pi_{j-1} = \mathrm{Exc}(f,\pi_1\circ\cdots\circ\pi_{j-1})\, u_2^{k_2}\cdots u_j^{k_j} x_j^r y_j^s.$$

Using Notation 1.21 and Lemma 1.22 we deduce that:

$$(\mathcal{M}_j(r,s), f)_0 = \mathrm{ord}_t(\mathrm{Exc}(f,\pi_1\circ\cdots\circ\pi_{j-1})) + r\,\mathrm{ord}_t(x_j) + s\,\mathrm{ord}_t(y_j) = e_{j-2}\bar{b}_{j-1} + r e_{j-1} + s\,(\bar{b}_{j+1} - n_j\bar{b}_j) \quad \square.$$

Lemma 2.7. *If $0 \leq r$ and $0 \leq s < e_{j-1}$ the Newton polygon of a term $\mathcal{M}_j(r,s)$, with respect to the coordinates (x_1, y_1), is contained in $\mathcal{N}(f)$ for $2 \leq j \leq g+1$. It is contained in the interior of $\mathcal{N}(f)$ unless $j = 2$, $r = 0$ and $0 \leq s < e_1$.*

Proof. We suppose that $0 \leq s < e_{j-1}$. If $j = 2$ we have that $\mathcal{M}_2(r,s) = x_1^{i_0} y_1^{i_1} f_2^s$ by Lemma 2.3. The vector $\vec{v} := (i_0 + s m_1, i_1)$ is a vertex of the Newton polygon of $\mathcal{M}_2(r,s)$ and $\vec{w} := (b_1, 0)$ is a vertex of $\mathcal{N}(f)$. The vector $\vec{p}_1 = (n_1, m_1)$ is orthogonal to the unique compact edge of the polyhedron $\mathcal{N}(f)$ and also of $\mathcal{N}(\mathcal{M}_1(r,s))$. We deduce the inequality:

$$n_1 b_1 = \langle \vec{p}_1, \vec{w}\rangle \leq \langle \vec{p}_1, \vec{v}\rangle = n_1 i_0 + s n_1 m_1 + m_1 i_1 = e_1 n_1 m_1 + r(m_1 c_1 - n_1 d_1) \overset{(15)}{=} n_1 b_1 + r,$$

from a simple calculation using (25). Equality holds in formula above if and only if $r = 0$.

If $j > 2$ we follow the proof of Proposition 2.1: there exist integers $0 < i_1 \leq n_1$, $0 < i'_0, k_2$ such that (27) holds. By Lemma 2.2 we have that $k_2 = l + [\frac{c_1 i'_0}{n_1}]$ where the integer l is $l := e_1 - i_2 - n_2 i_3 - \cdots - n_2 \cdots n_{j-1} i_j$. The vector $\vec{v} := (i_0 + m_1(e_1 - l), i_1)$ is a vertex of $\mathcal{N}(\mathcal{M}_j(r,s))$. By construction the Newton polygon of $\mathcal{M}_j(r,s)$ has at most one compact edge parallel to Γ_1. We deduce from a simple calculation using (24) that:

$$n_1 b_1 \leq \langle \vec{p}_1, \vec{v}\rangle = n_1 b_1 + i'_0. \tag{31}$$

By the proof of Proposition 2.1 we have that $i'_0 > 0$, hence the inequality (31) is strict. □

Remark 2.8. *By induction using the same arguments as in Lemma 2.7 we check that if $1 \leq i < j$, $0 \leq r$, and $0 \leq s < e_{j-1}$ that the Newton polygon of*

$$\mathcal{M}_j(r,s)\circ\pi_1\circ\cdots\circ\pi_{i-1}/Exc(f,\pi_1\circ\cdots\circ\pi_{i-1})$$

with respect to the coordinates (x_i, y_i), is contained in $\mathcal{N}(f^{(i)})$. It is contained in the interior of $\mathcal{N}(f^{(i)})$ unless $j = i+1$, $r = 0$ and $0 \leq s < e_i$.

3. Multi-semi-quasi-homogeneous deformations

In this Section we introduce a class of (non equisingular) deformations of a plane branch $(C, 0)$ and some basic properties. These deformations, which we call multi-semi-quasi-homogeneous (msqh), are applied in [GP-R] to describe smoothings with the maximal number of connected components.

We suppose that $(C, 0)$ is a plane branch. We keep notations of Section 1.4 and Notation 3.1. The resolution is described in terms of a fixed sequence $f_1, \ldots, f_g$ of normalized semi-roots (see Definition 1.14). We introduce the following notations:

Notation 3.1. *For $j = 1, \ldots, g$ we set:*

(i) $\Gamma_j = [(m_j e_j, 0), (0, n_j e_j)]$ *the unique compact edge of the local Newton polygon of $f^{(j)}(x_j, y_j)$ (see (19)),*

(ii) Δ_j *the triangle bounded by the Newton polygon of $f^{(j)}(x_j, y_j)$ and the coordinate axis; we denote by Δ_j^- the set $\Delta_j^- = \Delta_j \setminus \Gamma_j$,*

(iii) $\Xi_j = [(0,0), (0, n_j e_j)]$ *the edge of Δ_j which intersects the vertical axis,*

(iv) *Let $\omega_j : \Delta_j \cap \mathbf{Z}^2 \to \mathbf{Z}$ be defined by*

$$\omega_j(r,s) = e_j(e_j n_j m_j - r n_j - s m_j). \tag{32}$$



FIGURE 1. The Newton diagrams in Notation 3.1

We denote $t_j, \dots, t_g$ by $\underline{t}_j$, for any $1 \leq j \leq g$. We abuse of notation by denoting f by $P_{\underline{t}_{g+1}}$, for instance, in the following algebraic expressions:

$$\left\{ \begin{array}{lllllll} P_{\underline{t}_g} & := & P_{\underline{t}_{g+1}} & + & \displaystyle\sum_{(r,s) \in \Delta_g^- \cap \mathbf{Z}^2} A_{r,s}^{(g)} & t_g^{\omega_g(r,s)} & \mathcal{M}_g(r,s) \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ P_{\underline{t}_1} & := & P_{\underline{t}_2} & + & \displaystyle\sum_{(r,s) \in \Delta_1^- \cap \mathbf{Z}^2} A_{r,s}^{(1)} & t_1^{\omega_1(r,s)} & \mathcal{M}_1(r,s). \end{array} \right. \tag{33}$$

The terms $A_{r,s}^{(j)}$ are some constants while the t_j are considered as parameters, for $j = 1, \dots, g$. We suppose that $A_{0,0}^{(j)} \neq 0$.

Definition 3.2. *We denote by $C_{\underline{t}_l}$, or by $C_{\underline{t}_l}^{(1)}$, the deformation of $(C, 0)$ defined by $P_{\underline{t}_l}$, for $0 < t_1 \ll \cdots \ll t_l \ll 1$ and $l = 1, \dots, g$. We say that such a deformation is* multi-semi-quasi-homogeneous *(see* [GP-R]*).*

We denote by $C_{l,\underline{t}}^{(j)} \subset Z_j$ the strict transform of $C_{l,\underline{t}}$ by the composition of toric maps $\Pi_{j-1} \circ \cdots \circ \Pi_1$ and by $P_{\underline{t}_l}^{(j)}(x_j, y_j)$ (resp. by $\bar{P}_{\underline{t}_l}^{(j)}(x_j, u_j)$) the polynomial defining $C_{\underline{t}_l}^{(j)}$ in the coordinates (x_j, y_j) (resp. (x_j, u_j)), for $2 \leq j \leq l \leq g$. These notations are analogous to those used for C in Section 1.4, see (17). In particular, we have that the result of substituting in $P_{\underline{t}_j}^{(j)}(x_j, y_j)$, the term y_j by using (18), is $\bar{P}_{\underline{t}_l}^{(j)}(x_j, u_j)$. The deformation $C_{\underline{t}_j}^{(j)}$ is a *semi-quasi-homogeneous deformation* of $C_{\underline{t}_{j+1}}^{(j)}$ (see [GP-R] for the definition and more details). This motivates the choice of terminology. Notice that by the discussion of Section 1.3 the msqh-deformations can be embedded naturally in the same space as the monomial curve associated to $(C, 0)$.

Proposition 3.3. *If $1 \leq j < l \leq g$ the curves $C_{\underline{t}_l}^{(j)}$ and $C^{(j)}$ meet the exceptional divisor of $\Pi_{j-1} \circ \cdots \circ \Pi_1$ only at the point $o_j \in E_j$ and with the same intersection multiplicity e_{j-1}.*

Proof. If $j = 1$ we have that f and $P_{\underline{t}_l}$ have the same Newton polygon and moreover the symbolic restrictions of these two polynomials to the compact face Γ_1 of the Newton polygon coincide by Lemma 2.7. If $j > 1$ we show the result by induction using Remark 2.8. □

Proposition 3.4. *If $1 < j \leq g$ then E_j is the only irreducible component of the exceptional divisor of $\Pi_{j-1} \circ \cdots \circ \Pi_1$ which intersects $C_{\underline{t}_j}^{(j)}$ at e_{j-1} points counted with multiplicity if $0 < |t_j| \ll 1$. More precisely, we have that:*

(i) *The symbolic restriction of $P_{\underline{t}_{j+1}}^{(j)}(x_j, y_j)$ to the edge Γ_j of its local Newton polygon is of the form: $\alpha_j \prod_{s=1}^{e_j} \left(y_j^{n_j} - (1 + \gamma_s^{(j)} t_{j+1}^{e_{j+1} m_{j+1}}) x_j^{m_j} \right)$, with $\alpha_j, \gamma_s^{(j)} \in \mathbf{C}^*$, for $s = 1, \dots, e_j$.*

(ii) *The points of intersection of E_{j+1} with $C_{\underline{t}_{j+1}}^{(j+1)}$ are those with coordinates $x_{j+1} = 0$ and*

$$u_{j+1} = (1 + \gamma_s^{(j)} t_{j+1}^{e_{j+1} m_{j+1}})^{-1}, \text{ for } s = 1, \dots, e_j. \tag{34}$$

Proof. If $j = 1$ we have that the terms of the expansion of $P_{\underline{t}_2}$ which have exponents on the compact face of the Newton polygon $\mathcal{N}(f)$ are f and $\mathcal{M}_2(0, s)$ for $0 \leq s < e_1$ by Lemma 2.7. By Proposition 2.1 we have that $\mathcal{M}_2(0, s) = x_1^{m_1(e_1-s)} f_2^s$. The restriction of the polynomial $f + \sum_{s=0}^{e_1-1} A_{(0,s)}^{(2)} t_2^{e_2 m_2 (e_1-s)} x_1^{m_1(e_1-s)} f_2^s$ to the face Γ_j is equal to:

$$(y_1^{n_1} - x_1^{m_1})^{e_1} + \sum_{s=0}^{e_1-1} A_{(0,s)}^{(2)} (t_2^{e_2 m_2} x_1^{m_1})^{e_1-s} (y_1 - x_1^{m_1})^s. \tag{35}$$

Let us consider the polynomial $Q_1(V_1, V_2) := V_1^{e_1} + \sum_{s=0}^{e_1-1} A_{(0,s)}^{(2)} V_1^s V_2^{e_1-s}$. By hypothesis $A_{(0,0)}^{(2)} \neq 0$ hence the homogeneous polynomial Q_1 factors as: $Q_1(V_1, V_2) = \prod_{s=1}^{e_1} (V_1 - \gamma_s^{(1)} V_2)$ for some $\gamma_s^{(1)} \in \mathbf{C}^*$. The expression (35) is of the form: $Q_1(y_1^{n_1} - x_1^{m_1}, t_2^{m_2 e_2} x_1^{m_1}) = \prod_{s=1}^{e_1} (y^{n_1} - (1 + \gamma_s^{(1)} t_2^{m_2 e_2}) x_1^{m_1})$. This proves the first assertion in this case. The second follows from this by the discussion of Section 1.4.

If $j > 1$ we deduce by induction, following the proof of Proposition 2.1 and using Remark 2.8, that the restriction of $P_{\underline{t}_{j+1}}^{(j-1)}$ to the compact face of $\mathcal{N}(f^{(j-1)})$ is of the form: $(y_j^{n_j} - x_1^{m_j})^{e_j} + \sum_{s=0}^{e_j-1} A_{(0,s)}^{(j+1)} (t_{j+1}^{e_{j+1} m_{j+1}} x_j^{m_{j+1}})^{e_j-s} (y_j - x_j^{m_j})^s$. The result in this case follows by the same argument. □.

3.1. **Milnor number and generic msqh-smoothings.** If $(D, 0) \subset (\mathbf{C}^2, 0)$ is the germ of a plane curve singularity, defined by $h = 0$, for $h \in \mathbf{C}\{x, y\}$ reduced, we denote by $\mu(h)_0$ or by $\mu(D)_0$ the *Milnor number*, $\mu(h)_0 := \dim_{\mathbf{C}} \mathbf{C}\{x, y\}/(f_x, f_y)$. We have the following formula (see [R] and [Z2]):

$$\mu(h)_0 = (h, \frac{\partial h}{\partial y})_0 - (h, x)_0 + 1. \tag{36}$$

The Milnor number of the plane branch $(C, 0)$ expresses in terms of the generators of the semigroup Λ with respect to the coordinate line $x = 0$, by using the classical formula (see [M], [GB] and [Z2]):

$$(f, \frac{\partial h}{\partial y})_0 = \sum_{j=1}^{g} (n_j - 1) \bar{b}_j, \tag{37}$$

Definition 3.5. *We say that the deformation $P_{\underline{t}_1}$ of the plane branch $(C, 0)$ is* generic *if the numbers $\{\gamma_s^{(j)}\}_{s=1}^{e_j}$ appearing in Proposition 3.4 are all distinct, for $1 < j \leq g$.*

Proposition 3.6. *Let $P_{\underline{t}_1}$ be a generic msqh-deformation of a plane branch $(C, 0)$, then we have that $\mu(C)_0 = \displaystyle\sum_{j=1}^{g} (\mu(C_{\underline{t}_{j+1}}^{(j)})_{o_j} + e_j - 1)$.*

Proof. We prove the result by induction on g. If $g = 1$ the assertion is trivial. We suppose the assertion true for branches with $g - 1$ characteristic exponents with respect to some system of coordinates. By Proposition 1.18 and the induction hypothesis it is easy to check that $\mu(C^{(2)})_{o_2} = \sum_{j=2}^{g} (\mu(C_{\underline{t}_{j+1}}^{(j)})_{o_j} + e_j - 1)$.

By Proposition 3.4 and the definition of generic msqh-deformation, we have that the curve $C_{\underline{t}_2}^{(1)}$, defined by the polynomial $P_{\underline{t}_2}^{(1)}(x_1, y_1)$, is non degenerated with respect to its Newton polygon, for $0 < |t_2| \ll 1$. By (36) we have that $\mu(C_{\underline{t}_2}^{(1)})_{o_1} = e_0 b_1 - e_0 - b_1 + 1$.

By (37) we have that: $\mu(C)_{o_1} - \mu(C^{(2)})_{o_2} = (f, \frac{\partial f}{\partial y})_0 - (f^{(2)}, \frac{\partial f^{(2)}}{\partial y_2})_{o_2} + e_1 - e_0$. The assertion holds if and only if $(f, \frac{\partial f}{\partial y})_0 - (f^{(2)}, \frac{\partial f^{(2)}}{\partial y_2})_{o_2} = b_1(e_0 - 1)$. Using (37) and Lemma 1.22 we verify that $(f, \frac{\partial f}{\partial y})_0 - (f^{(2)}, \frac{\partial f^{(2)}}{\partial y_2})_{o_2}$ is equal to:

$$\begin{array}{rcl} \sum_{j=1}^{g}(n_k-1)\bar{b}_j - \sum_{j=2}^{g}(n_k-1)\left(\bar{b}_j - \bar{b}_1 e_0/e_{j-1}\right) & = & (n_1-1)\bar{b}_1 + \sum_{j=2}^{g}\bar{b}_1 e_0/e_{j-1} = \\ (n_1 - 1 + (n_2-1)n_1 + \cdots + (n_g-1)n_1\cdots n_{g-1})\,\bar{b}_1 & = & (e_0-1)b_1. \quad \square \end{array}$$

Definition 3.7. *We associate to the plane branch $(C,0)$ an abstract simplicial complex $\Omega(C)$ with integral structure, by gluing the edge Γ_{j-1} of the triangle Δ_{j-1} with the edge Ξ_j of Δ_j, for $j = 1,\ldots,g-1$. The integral structure is preserved since the edges Γ_{j-1} and Ξ_j have the same integral length e_j. If $g=1$ then $\Omega(C) = \Delta_0$ and this complex is embedded in $\mathbf{R}^2$.*

The following consequence of Proposition 3.6 generalizes Kouchnirenko's formula (see [K]).

Proposition 3.8. *The Milnor number of the plane branch $(C,0)$ is twice the number of integral points in the simplicial complex with integral structure $\Omega(C)$. Namely, we have that:*

$$\mu(C)_0 = 2\sum_{j=0}^{g-1}\left(\#(\overset{\circ}{\Delta}_j \cap \mathbf{Z}^2) + e_{j+1} - 1\right).$$

See [G] for related formulas, measuring the difference of the Milnor number or the delta invariant, of a reduced plane curve germ and its strict transform by a suitable toric modification.

4. IRREDUCIBILITY AND EQUISINGULARITY CRITERIONS

Abhyankar's irreducibility criterion gives an affirmative answer to a question of Kuo mentioned in [Abh4]: "Can we decide the irreducibility of a power series $F(x,y)$ without blowing up and without getting into fractional power series ?" We have revisited the Abhyankar's criterion in the light of toric geometry methods. In particular, our proof explains that if F is irreducible, some information on Newton polygons of the strict transform of F at the infinitely near points of the toric resolution can be read from the expansions in certain approximate roots of F. See [C-M2] and [C-M1], for an extension of this criterion to the case of base field of positive characteristic. Recently, García Barroso y Gwoździewicz have given another method to answer Kou's question based on the analysis of the jacobian Newton polygons of plane branches (see [GB-G], see also Section 4.3.1)

As an application we give an equisingularity criterion for an equimultiple family of plane curves to be equisingular to a plane branch (See Section 4.3).

4.1. **Straight line conditions in the toric resolution.** We study the compatibility of the expansions in terms of semi-roots with the toric resolution.

In this section, we set f_j Weierstrass polynomial of degree $n_0 \ldots n_{j-1}$, defining a semi-root of f, for $j = 1,\ldots,g$. We give further precisions of the $(y = f_1,\cdots,f_j)$-expansion of f (see Proposition 1.9).

Proposition 4.1. *We have that the $(f_1,\cdots,f_j)$-expansion of f, for $2 \le j \le g$, is of the form:*

$$f = f_j^{e_{j-1}} + \sum_{I=(i_1,\ldots,i_j)} \alpha_I(x_1) f_1^{i_1}\cdots f_j^{i_j}, \text{ for } \alpha_I(x_1) \in \mathbf{C}\{x_1\}, \tag{38}$$

with $0 \le i_1 < n_1$, ..., $0 \le i_{j-1} < n_{j-1}$, $0 \le i_j < e_{j-1}$, where:

1. *The orders in y_1 of the series $(f_1^{i_1}\cdots f_j^{i_j})_{|x_1=0}$, corresponding to the terms $f_1^{i_1}\cdots f_j^{i_j}$ in expansion (38), are all distinct.*

(2) *The Newton polygons of* f *and of* $f_j^{e_{j-1}}$*, with respect to the coordinates* (x_1, y_1) *have only one compact edge* Γ_1*, defined in section 1.4, and the symbolic restrictions of* f *and of* $f_j^{e_{j-1}}$ *coincide on this edge.*

(3) *If* $\alpha_I \neq 0$ *the Newton polygon of the term* $\alpha_I(x_1) f_1^{i_1} \cdots f_j^{i_j}$ *is contained in the interior of the Newton polygon of* f*.*

Proof. Since $\deg f = e_{j-1} \deg f_j$ and both are monic polynomials we have that the term $f_j^{e_{j-1}}$ appears in the $(f_1, \cdots, f_j)$-expansion of f with coefficient one.

Let $\mathcal{M}_I := \alpha_I(x) f_1^{i_1} \cdots f_j^{i_j}$ be a term of the expansion (38) appearing with non zero coefficient $\alpha_I(x)$. By Proposition 1.9 we have that $0 \leq i_1 < n_1, \ldots, 0 \leq i_{j-1} < n_{j-1}$ and $0 \leq i_j < e_{j-1}$. The assertion on the orders in y_1 of the series $(f_1^{i_1} \cdots f_j^{i_j})_{|x_1=0}$, follows from Remark 1.10. By Remark 1.17, the term $\mathcal{M}_I$ has at most one compact edge Γ_I, which is parallel to the compact face Γ_1 of $\mathcal{N}(f)$. The vector $\vec{p_1} = (n_1, m_1)$, which was defined in Section 1.4, is orthogonal to the compact face Γ_1 of $\mathcal{N}(f)(x_1, y_1)$. We set $c_I := \min\{\langle \vec{p_1}, \vec{u} \rangle \mid \vec{u} \in \mathcal{N}(\mathcal{M}_I)\}$. Suppose that there exists a term $\mathcal{M}_I$, with $\alpha_I \neq 0$ such that: $c_I < \min\{\langle \vec{p_1}, \vec{u} \rangle \mid \vec{u} \in \mathcal{N}(f)\}$. In that case we can choose an index I, with minimal c_I and such that: $\mathrm{ord}_{y_1}(f_1^{i_1} \cdots f_j^{i_j})_{|x_1=0} = \max\{\mathrm{ord}_{y_1}(f_1^{i'_1} \cdots f_j^{i'_j})_{|x_1=0} \mid c_I = c_{I'}\}$. Notice that this index I is unique since the orders, $\mathrm{ord}_{y_1}(f_1^{i_1} \cdots f_j^{i_j})_{|x_1=0}$, are all distinct. We have that if $(r, s) \in \Gamma_I \cap \mathbf{Z}^2$, the sum K_I of the coefficients of the term $x^r y^s$ in $\alpha_I \mathcal{M}'_I$, for those I' with $c_I = c_{I'}$ must vanish. But if (r, s) is the vertex of Γ_I with $s = \deg_y f_1^{i_1} \cdots f_j^{i_j}$ then we obtain that K_I is the initial coefficient of the series α_I, a contradiction.

By (14) the symbolic restrictions of f and of $f_j^{e_{j-1}}$, to the compact face Γ_1 of the Newton polygon coincide. This implies that if there exists a term $\mathcal{M}_I$ with $c_I = \min\{\langle \vec{p_1}, u \rangle \mid u \in \mathcal{N}(f)\}$, then the sum of the coefficients of the terms $x^r y^s$ in $\mathcal{M}'_I$, for those I' with $c_I = c_{I'}$, must vanish. We argue as in the previous case, to prove that cannot happen. □

Remark 4.2. *Notice that* f_j *is an approximate root of* f *if and only if the exponents* i_j*, appearing in (38), verify that* $i_j < e_{j-1} - 1$*.*

The $(x_1, f_1, \ldots, f_j)$-expansion of f

$$f = f_j^{e_{j-1}} + \sum_{I=(i_0,\ldots,i_j)} c_I \, x_1^{i_0} \, f_1^{i_1} \cdots f_j^{i_j}, \tag{39}$$

is obtained by expanding, as a series in x_1, the coefficients of the terms in (38). If $c_I \neq 0$, then the Newton polygon of the term $\mathcal{M}_I := x^{i_0} f_1^{i_1} \cdots f_j^{i_j}$ is contained in the interior of the Newton polygon of f by Proposition 4.1. It follows from the discussion in Section 1.4 that we can factor $\mathcal{M}_I \circ \pi_1$ as:

$$\mathcal{M}_I \circ \pi_1 = \mathrm{Exc}(f, \pi_1) \, u_2^{k_2(I)} \, x_2^{i'_0(I)} \, (f_2^{(2)})^{i_2} \cdots (f_j^{(2)})^{i_j}, \tag{40}$$

where $k_2(I)$ and $i'_0(I)$ are non negative integers. More precisely, if we set $q_I := i_1 + n_1 i_2 + \cdots + n_1 \cdots n_{j-1} i_j$, then we have that $k_2(I) = c_1 i_0 - d_1(e_0 - q_I) \geq 0$ and $i'_0(I) = n_1 i_0 - m_1(e_0 - q_I) \geq 0$.

We obtain the following expansion from (39), by factoring out $\mathrm{Exc}(f, \pi_1)$ from $f \circ \pi_1$:

$$f^{(2)} = (f_j^{(2)})^{e_{j-1}} + \sum_{I=(i_0,\ldots,i_j)} c_I \, u_2^{k_2(I)} \, x_2^{i'_0(I)} \, (f_2^{(2)})^{i_2} \cdots (f_j^{(2)})^{i_j}. \tag{41}$$

The following expansion is obtained from (41) by collecting the terms with the same index $I' = (i_2, \ldots, i_j)$:

$$f^{(2)} = (f_j^{(2)})^{e_{j-1}} + \sum_{I'=(i_2,\ldots,i_j)} \alpha_{I'}^{(2)}(x_2, u_2) \, (f_2^{(2)})^{i_2} \cdots (f_j^{(2)})^{i_j}. \tag{42}$$

By (18) the coefficient $\alpha_{I'}^{(2)}(x_2, u_2)$, viewed in $\mathbf{C}\{x_2, y_2\}$, is of the form:

$$\alpha_{I'}^{(2)} = \epsilon_{I'}^{(2)} x_2^{r_2(I')} \text{ with } r_2(I') \geq 0 \text{ and } \epsilon_{I'}^{(2)} \text{ a unit in } \mathbf{C}\{x_2, y_2\}. \tag{43}$$

Definition 4.3. *We call the expansion (41) (resp. (42)) the* $(u_2, x_2, f_2^{(2)}, \dots, f_j^{(2)})$*-expansion (resp.* $(f_2^{(2)}, \dots, f_j^{(2)})$*-expansion) of* $f^{(2)}$*.*

Proposition 4.4. *Suppose that* $2 \leq j \leq g$*. The expansion (42) verifies that:*

1. *The orders in* $y_2 = f_2^{(2)}$ *of the series* $((f_2^{(2)})^{i_2} \cdots (f_j^{(2)})^{i_j})_{|x_2=0}$*, corresponding to the terms* $(f_2^{(2)})^{i_2} \cdots (f_j^{(2)})^{i_j}$ *in expansion (42), are all distinct.*
2. *If* $g > 2$*, the Newton polygons of* $f^{(2)}$ *and of* $(f_j^{(2)})^{e_{j-1}}$*, with respect to the coordinates* (x_2, y_2) *have only one compact edge* Γ_2 *(see section 1.4). Moreover, then the symbolic restrictions of* $f^{(2)}$ *and of* $(f_j^{(2)})^{e_{j-1}}$ *coincide on this edge.*
3. *If* $g > 2$ *and if* $\alpha_{I'}^{(2)}(x_2, u_2) \neq 0$*, then the Newton polygon of the term:*
$$\alpha_{I'}^{(2)}(x_2, u_2)\, (f_2^{(2)})^{i_2} \cdots (f_j^{(2)})^{i_j}$$
is contained in the interior of $\mathcal{N}(f^{(2)})$*, with respect to the coordinates* (x_2, y_2)*.*
4. *If* $j = g = 2$*, the term of the expansion (42) corresponding to* $i_2 = 0$ *is of the form* $x_2^{\bar{b}_2 - n_1 \bar{b}_1} \epsilon_0^{(2)}$*, where* $\epsilon_0^{(2)}$ *is a unit in* $\mathbf{C}\{x_2, y_2\}$*. The terms of the expansion (42) different from* $y_2^{e_1} = (f_2^{(2)})^{e_1}$ *and* $x_2^{\bar{b}_2 - n_1 \bar{b}_1} \epsilon_0^{(2)}$*, have Newton polygon contained in the interior of* $\mathcal{N}(f^{(2)})$*.*

Proof. By Proposition 1.18 the characteristic integers of $(C_j^{(2)}, o_2)$, with respect to the line $x_2 = 0$ are $n_2, \dots, n_{j-1}$. The assertion on the orders in y_2 of the series $((f_2^{(2)})^{i_2} \cdots (f_j^{(2)})^{i_j})_{|x_2=0}$ is consequence of Remark 1.10. Suppose first that $g > 2$. We have that if $2 < j \leq g$, the term $f_j^{(2)}$ is of the form $f_j^{(2)} = (y_2^{n_2} - \theta_2 x_2^{m_2})^{n_3 \dots n_{j-1}} + \cdots$, where the terms which are not written in this expansion have exponents contained in the interior of the Newton polygon with respect to the coordinates x_2 and y_2. Namely, by Remark 1.17 and the discussion of Section 1.4 we have that $f^{(2)}$ and of $(f_j^{(2)})^{e_{j-1}}$, have the same Newton polygon with respect to the coordinates (x_2, y_2). This polygon has only one edge Γ_2. The symbolic restrictions of $f^{(2)}$ and of $(f_j^{(2)})^{e_{j-1}}$ coincide on this edge by Remark 1.17 and Proposition 1.18.

It follows that the Newton polygon of a term $\mathcal{M}_{I'}^{(2)} := \alpha_{I'}^{(2)}(x_2, u_2)\, (f_2^{(2)})^{i_2} \cdots (f_j^{(2)})^{i_j}$, appearing in the expansion (42), has only one compact face which is parallel to Γ_2. We deduce that the Newton polygon of $\mathcal{M}_I^{(2)}$ is contained in the interior of $\mathcal{N}(f)$ by repeating the argument of Proposition 4.1.

If $j = g = 2$ then it is clear that the Newton polygon of $f^{(2)}$ with respect to the coordinates (x_2, y_2) is equal to the convex hull of the set $(0, e_1) + \mathbf{R}_{\geq 0}^2 \cup \bigcup_{\alpha_{I'}^{(2)} \neq 0} (r_2, i_2) + \mathbf{R}_{\geq 0}^2$. By Proposition 1.18 we know that this Newton polygon has only one compact edge Γ_2 with no integral point other than the vertices $(0, e_1)$ and $(\bar{b}_2 - n_1 \bar{b}_1, 0)$. □

Theorem 4.1. *The* $(x, f_1, \dots, f_j)$*-expansion of* f*, for* $j = 2, \dots, g$*, is of the form:*

$$f = f_j^{e_{j-1}} + \sum c_I\, \mathcal{M}_j(r(I), s(I)),$$

where $(r(I), s(I))$ *verify that* $0 < r(I)$*,* $0 \leq s(I) < e_{j-1}$*,* $r(I) e_{j-1} + s(I)(\bar{b}_j - n_{j-1}\bar{b}_{j-1}) \geq e_{j-1}(\bar{b}_j - n_{j-1}\bar{b}_{j-1})$*. Among the terms of this expansion with minimal intersection multiplicity with* f *there exist* $f_j^{e_{j-1}}$ *and* $\mathcal{M}_j(\bar{b}_{j+1} - n_j \bar{b}_j, 0)$*. Moreover, if* $j = g - 1$ *these two terms are exactly the terms with minimal intersection multiplicity with* f*.*

Let $\mathcal{M}_I := x^{i_0} f_1^{i_1} \cdots f_j^{i_j}$ be a monomial appearing in (39). Using Proposition 4.1 and Proposition 4.4, we deduce inductively that we can factor $\mathcal{M}_I \circ \pi_1 \circ \cdots \circ \pi_{j-1}$ as

$$\mathcal{M}_I \circ \pi_1 \circ \cdots \circ \pi_{j-1} = \operatorname{Exc}(f, \pi_1 \circ \cdots \circ \pi_{j-1}) u_2^{k_2(I)} \cdots u_j^{k_j(I)} x_j^{r(I)} y_j^{s(I)}, \tag{44}$$

where $k_2(I), \ldots, k_j(I), r(I) \geq 0$ and $s(I) = i_j$. It follows that we have an expansion:

$$f^{(j)} = y_j^{e_{j-1}} + \sum_I c_I \, u_2^{k_2(I)} \cdots u_j^{k_j(I)} \, x_j^{r(I)} \, y_j^{s(I)}. \tag{45}$$

By (18) the term $u_2^{k_2(I)} \cdots u_j^{k_j(I)}$ is a unit viewed in $\mathbf{C}\{x_j, y_j\}$.

If $I \neq I'$ are two different indices appearing in (39), then $(r(I), s(I)) \neq (r(I'), s(I'))$ by Proposition 2.1. Hence, the Newton polygon of $f^{(j)}(x_j, y_j)$ is equal to the convex hull of the set, $\cup_I (r(I), s(I)) + \mathbf{R}^2_{\geq 0}$. By Proposition 1.18 at level j, we have that this polygon has vertices $(0, e_{j-1})$ and $(\bar{b}_j - n_{j-1}\bar{b}_{j-1}, 0)$. If $j = g$, these two vertices are the unique integral points in the Newton polygon. By Lemma 2.6, the exponents $(r, s) \in \Gamma_j$ correspond to terms $\mathcal{M}_j(r, s)$ with minimal intersection multiplicity with f at the origin. □

4.2. **Abyankar's generalized Newton polygons, straight line condition and irreducibility criterion.** We follow the presentation given by Assi and Barile in [As-B] of results in [Abh4].

4.2.1. *Generalized Newton polygons.* Given a sequence $\underline{\bar{B}} := (\bar{B}_0, \bar{B}_1 < \cdots < \bar{B}_G)$ of positive integers, we associate to them sequences $E_j = \gcd(\bar{B}_0 = N, \bar{B}_1, \ldots, \bar{B}_G)$ and $N_0 = 1$, $N_j = E_{j-1}/E_j$, for $j = 0, \ldots, G$. Notice that if $\underline{\bar{B}}$ a characteristic sequence of generators of the semigroup Λ associated to a plane branch $(C, 0)$, we set $g = G$ and we have with the Notations of the first section that $E_j = e_j$ and $N_j = n_j$, for $j = 0, \ldots, g$.

Let

$$F = y^N + \sum_{i=2}^{N} A_i(x) y^{n-i} \in \mathbf{C}\{x\}[y] \tag{46}$$

be a Weierstrass polynomial. We assume that y is an approximate root of F, since the coefficient of y^{N-1} is equal to zero. We denote by F_j the approximate root of F of degree $N_0 \ldots N_{j-1}$, and by $\underline{F}_j$ the sequence $(F_1, \ldots, F_j)$ for $j = 1, \ldots, G+1$ and $\underline{F} = \underline{F}_{g+1}$.

Let $P \in \mathbf{C}\{x\}[y]$ be a monic polynomial. We consider the $(F_1, \ldots, F_{G+1})$-expansion $P = \sum_I \alpha_I(x) F_1^{i_1} \cdots F_G^{i_G} F_{g+1}^{i_{G+1}}$ of P (see Proposition 1.9). The *formal intersection multiplicity* of P and $\underline{F}$, with respect to the sequence $\underline{\bar{B}}$ is defined as

$$\text{formal}_{\,\underline{\bar{B}}}(P, \underline{F}) := \min\{\sum_{j=0}^{G} i_j \bar{B}_j \mid I = (i_1, \ldots, i_G, 0), \alpha_I(x) \neq 0\}. \tag{47}$$

Notice that when this value is $< +\infty$, it is reached at only one coefficient.

Let $P, Q \in \mathbf{C}\{x\}[y]$ be two monic polynomials of degrees p, q with $p = mq$. We have the Q-adic expansion of P is of the form: $P = Q^m + \alpha_1 Q^{m-1} + \cdots + \alpha_m$. The *generalized Newton polygon* of P with respect to Q and the sequences $\underline{\bar{B}}$ and $\underline{\bar{F}}$ is the convex hull of the set:

$$\mathcal{N}(P, Q, \underline{\bar{B}}, \underline{F}) := \bigcup_{k=0}^{m} (\text{formal}_{\,\underline{\bar{B}}}(\alpha_k, \underline{F}), (d-k)\text{formal}_{\,\underline{\bar{B}}}(Q, \underline{F})) + \mathbf{R}^2_{\geq 0}. \tag{48}$$

4.2.2. *Abhyankar's irreducibility criterion.* To a monic polynomial F of the form (46) it is associated a sequence $\bar{B}$ as follows: Set $\bar{B}_0 = E_0 := N$, $F_1 = Y$, $\bar{B}_1 = (F_1, F)_0$, $E_1 = \gcd(\bar{B}_0, \bar{B}_1)$ and $N_1 := E_0/E_1$. Then, for $j \geq 2$ the integers $E_0 \ldots, E_j = \gcd(\bar{B}_0, \ldots, \bar{B}_j)$, and $N_1, \ldots, N_{j-1}$ are defined by induction. We set $B_k = (P, F_j)_0$ where F_j denotes the approximate root of F, of degree $N_1 \cdots N_{j-1}$.

Theorem 4.2. ([Abh4]) *With the above notations the polynomial $F \in \mathbf{C}\{x\}[y]$ is irreducible if and only if the following conditions hold:*

1. *There exists an integer $G \in \mathbf{Z}_{>0}$ such that $E_G = 1$.*
2. *We have that $\bar{B}_{j+1} > N_k \bar{B}_j$ for $j = 1, \ldots G - 1$.*
3. (straight line condition) *The generalized Newton polygon $\mathcal{N}(F_{j+1}, F_j, \frac{1}{E_j}\underline{\bar{B}_j}, \underline{F}_j)$ has only one compact edge with vertices $(\frac{1}{E_j} N_j \bar{B}_j, 0)$ and $(0, \frac{1}{E_j} N_j \bar{B}_j)$.*

Proof. We prove first that if F verifies the conditions of the theorem, then F is irreducible. By straight line condition the vertices of the generalized Newton polygon, $\mathcal{N}(F_{j+1}, F_j, \frac{1}{E_j}\underline{\bar{B}_j}, \underline{F}_j)$, correspond to the terms $F_j^{n_j}$ and $\alpha_0^{(j)}(x) F_1^{\eta_1^{(j)}} \cdots F_{j-1}^{\eta_{j-1}^{(j)}}$ of the $(F_1, \ldots, F_j)$ expansion of F_{j+1}, where $\mathrm{ord}_x \alpha_0^{(j)}(x) = \eta_0^{(0)} \geq 0$ and $0 \leq \eta_i < N_i$ for $i = 1, \ldots, j-1$. The straight line condition implies that $\frac{1}{E_j} N_j \bar{B}_j = \frac{1}{E_j}(\eta_0^{(j)} \bar{B}_0 + \cdots + \eta_{j-1}^{(j)} \bar{B}_{j-1})$. It follows that $N_j \bar{B}_j$ belongs to the semigroup generated by $\bar{B}_0, \ldots, \bar{B}_{j-1}$. By Lemma 1.2, this numerical condition together with the first two hypothesis of the theorem guarantee that the semigroup generated by $\bar{B}_0, \ldots, \bar{B}_g$ is the semigroup of a plane branch.

Let $0 \neq F' \in \mathbf{C}\{x\}[y]$ be any polynomial of degree $< N = \deg F$. We consider the $(F_1, \ldots, F_g)$ expansion of F':

$$F' = \sum_I \alpha_I(x) F_1^{i_1} \cdots F_g^{i_g} \text{ with } \alpha_I(x) \in \mathbf{C}\{x\}, \tag{49}$$

where the components of the index $I = (i_1, \ldots, i_g)$ verify that $0 \leq i_1 < N_1, \ldots, 0 \leq i_g \leq [\deg F'/\deg F_g] < n_g = \deg F/\deg F_g$. If $i_0 = \mathrm{ord}_x \alpha_I(x)$ we have that

$$(F, \alpha_I(x) F_1^{i_1} \cdots F_g^{i_g})_0 = \sum_{j=0}^{g} i_j \bar{B}_j. \tag{50}$$

It follows that from Lemma 1.2 that the intersection multiplicities (50) corresponding to different terms in the finite sum (49) are all different. We deduce that $(F, F')_0 = \min\{\sum_{j=0}^{g} i_j \bar{B}_j \mid \alpha_i(x) \neq 0\} < +\infty$. It follows that F is irreducible, otherwise there is an irreducible factor of F' of F of degree $< \deg F$ and then $(F, F')_0 = +\infty$, a contradiction.

Suppose now that F is irreducible. Then $\bar{B}_0, \ldots, \bar{B}_G$ are the generators of the semigroup of the branch $F = 0$ with respect to the line $x = 0$. By Lemma 1.2, the first two conditions in the statement of the theorem hold automatically. By Proposition 1.8 the approximate root F_{j+1} is irreducible and define a plane branch with semigroup generated by $\frac{1}{E_j}\bar{B}_0, \ldots, \frac{1}{E_j}\bar{B}_j$.

By Theorem 4.1 we have that the Newton polygon of $F_{j+1}^{(j)}(x_j, y_j)$ has only two vertices $(N_j, 0)$ and $(M_j, 0)$ which correspond to the terms $F_j^{N_j}$ and $\mathcal{M}_j^{(j+1)}(0, M_j)$ of the $(x, F_1, \ldots, F_j)$-expansion of F_{j+1}. By Proposition 1.18 and induction the vertices of the Newton polygon of the strict transform function $F^{(j)}(x_j, y_j)$ are $(\bar{B}_j - N_{j-1}\bar{B}_{j-1}, 0)$ and $(0, E_j N_j)$ (see Notations of Section 1.4). It follows that $M_j = \frac{1}{E_j}(\bar{B}_j - N_{j-1}\bar{B}_{j-1})$ By Lemma 1.22 we have that $\mathrm{ord}_t(\mathrm{Exc}(F_{j+1}, \pi_1 \circ \cdots \circ \pi_{j-1}) = \frac{E_{j-2}}{E_j}\frac{\bar{B}_{j-1}}{E_j} = \frac{1}{E_j} N_{j-1} N_j \bar{B}_j$. By Lemma 2.6 and 1.22

the number $(F_{j+1}, \mathcal{M}_j^{(j+1)}(0, M_j))_0$ is equal to:

$$\frac{1}{E_j} N_{j-1} N_j \bar{B}_{j-1} + \frac{E_{j-1}}{E_j} M_j = \frac{1}{E_j}(N_j N_{j-1} \bar{B}_{j-1} + N_j(\bar{B}_j - N_{j-1}\bar{B}_{j-1})) = \frac{1}{E_j} N_j \bar{B}_j = (F_{j+1}, F_j^{N_j})_0.$$

A similar computation using Lemma 2.6 and 1.22 proves that if $\mathcal{M}_j^{(j+1)}(r,s)$ appears in the $(x, F_1, \dots, F_j)$-expansion of F_{j+1} then $(F_{j+1}, \mathcal{M}_j^{(j+1)}(r,s))_0 > \frac{1}{E_j} N_j \bar{B}_j$. □

If F is irreducible certain generalized Newton polygons (48) express in terms of the Newton polygons appearing in the toric embedded resolution:

Proposition 4.5. *Let $\phi_j : \mathbf{R}^2 \to \mathbf{R}^2$ be the linear function given by $\phi_j(r,s) = (rN_j, sM_j)$. We denote by R_j the number $R_j := \frac{1}{E_j} N_{j-1} N_j \bar{B}_j$. If F is irreducible we have that:*

$$\mathcal{N}(F_{j+1}, F_j, \underline{\frac{1}{E_j}\bar{B}_j}, \underline{F}_j) = (R_j, R_j) + \phi_j(\mathcal{N}(F_{j+1}^{(j)}(x_j, y_j))).$$

4.3. **Equisingularity criterions.** Let $F = F(\lambda, x, y) \in \mathbf{C}\{\lambda, x, y\}$ such that $F(\lambda, 0, 0) = 0$. We consider a family $(C_\lambda, 0)$ of plane curve germs defined by the vanishing of $F_\lambda(x,y) := F(\lambda, x, y)$. As an application we give a criterion for the uniparametrical deformation C_λ to be an equisingular deformation at $\lambda = 0$ of a plane branch $(C, 0)$. Our criterion extends the one given by A'Campo and Oka in [A'C-Ok]. They assume that certain approximate roots of C_l do not depend on λ. We do not need this hypothesis.

We have an expansion of the form:

$$F_\lambda = (p(\lambda)x + q(\lambda)y)^N + \sum_{r+s>N} c_{r,s}(\lambda) x^r y^s. \tag{51}$$

The family $(C_\lambda, 0)$ is equimultiple at $\lambda = 0$ if and only if $p(0)$ and $q(0)$ do not vanish simultaneously. Without loss of generality we suppose that $q(0) \neq 0$. If we make a change of coordinates by setting $y' = p(\lambda)x + q(\lambda)y$ we reduce to the case when the coordinate line $y' = 0$ is tangent to $(C_\lambda, 0)$, for $|\lambda| \ll 1$. We reduce to the case of families of the form:

$$F_\lambda = y^N + \sum_{r+s>N} c_{r,s}(\lambda) x^r y^s. \tag{52}$$

Notice that $\mathrm{ord}_y F(0,0,y) = \mathrm{ord}_y F_\lambda(0,y)$ is equal to the multiplicity of $(C_\lambda, 0)$, for $|\lambda| \ll 1$. By the Weierstrass Preparation Theorem we can suppose that F is a Weierstrass polynomial in y, and also that y is an approximate root of F, by making a local change of coordinates. The polynomial F is of the form:

$$F = y^N + \sum_{i=2}^{N} A_{i,\lambda}(x) y^{n-i} \in \mathbf{C}\{x, \lambda\}[y]. \tag{53}$$

Our aim is to characterize when the family (C_λ, o_λ) is equisingular at $\lambda = 0$, with special fiber $(C_0, 0)$ being a plane branch (irreducible and reduced). Equisingular, in this case means that each fiber $(C_\lambda, 0)$ is analytically irreducible and reduced, with the same characteristic exponents with respect to the line $x = 0$. We suppose from now on that $\mathrm{mult}(C_0, 0) > 1$. Since y is the approximate root of degree 1 of F_λ we have that:

Proposition 4.6. *With the above notations and hypothesis if $(C_\lambda, 0)$ defines an equisingular family at $\lambda = 0$, then we have that F_λ is of the form:*

$$F_\lambda = (y_1^{n_1} - \theta_1(\lambda) x_1^{m_1})^{e_1} + \cdots, \quad \text{with } \theta_1(\lambda) \in \mathbf{C}\{\lambda\} \text{ and } \theta_1(0) \neq 0, \tag{54}$$

where gcd $(n_1, m_1) = 1$ *and the terms which are not written have exponents in the interior of the Newton polygon of F_λ.*

Proof. We have an inclusion of polygons $\mathcal{N}(F_0) \subset \mathcal{N}(F_\lambda)$. By hypothesis both polygons have a vertex at $(0, n)$. Since $(C_\lambda, 0)$ is analytically irreducible it follows that $\mathcal{N}(F_\lambda)$ has only one compact edge. Since the coefficient of y_1^{n-1} in F_λ is equal to zero, as a consequence of the Tschirnhausen transformation, the inclination of this edge is equal to the first characteristic exponent of $(C_\lambda, 0)$. We deduce that $\mathcal{N}(F_\lambda) = \mathcal{N}(F_0)$. If F_λ is not of the form (54), then $(C_\lambda, 0)$ is not be analytically irreducible. $\square$

Suppose that we know the sequence of characteristic exponents and integers associated to $(C_0, 0)$ with respect to the line $x = 0$. We use Notations of Section 1.1 for them. We denote the approximate root of degree $n_0 \cdots n_{j-1}$ by $F_{j,\lambda}$, and by $(C_{j,\lambda}, 0)$ the deformation defined by $F_{j,\lambda} = 0$, for $j = 1, \dots, g+1$. The polynomials $f_j := F_{j,0}$ are semi-roots for $(C_0, 0)$ with respect to $x = 0$, by Proposition 1.8.

The notation $\mathcal{M}_j^{(j+1)}(r, s)$, for $0 \leq r$ and $0 \leq s < n_j$, define monomials in terms of the semi-roots $x, f_1, \dots, f_j$ (see Definition 2.4). If $\mathcal{M}_j^{(j+1)}(r, s) = x^{i_0} f_1^{i_1} \cdots f_j^{i_j}$ we denote by $\mathcal{M}_{j,\lambda}^{(j+1)}(r, s)$ the term $\mathcal{M}_{j,\lambda}^{(j+1)}(r, s) := x^{i_0} F_{1,\lambda}^{i_1} \cdots F_{j,\lambda}^{i_j}$. We define the weight of the monomial $\mathcal{M} = \mathcal{M}_{j,\lambda}^{(j+1)}(r, s)$ as the sum $i_o \bar{b}_0 + \cdots i_j \bar{b}_j$.

We have the following result (compare with Proposition 1.13).

Theorem 4.3. *With the above hypothesis and notations if* $(C_\lambda, 0)$ *is equisingular at* $\lambda = 0$ *to the plane branch* $(C_0, 0)$ *if and only for* $j = 1, \dots, g$ *we have expansions of the form:*

$$F_{j+1,\lambda} = F_{j,\lambda}^{n_j} - \theta_{j+1}(\lambda)\, \mathcal{M}_{j,\lambda}^{(j+1)}(m_j, 0) + \sum \alpha_{r,s}^{(j+1)}(\lambda)\, \mathcal{M}_{j,\lambda}^{(j+1)}(r, s), \tag{55}$$

where $\theta_{j+1}(0) \neq 0$ *and the terms* $\mathcal{M}_{j,\lambda}^{(j+1)}(r, s)$*, which appear in the sum with non zero coefficient, have weights strictly bigger than* $n_j \bar{b}_j$*.*

Proof. If we have expansions of the form (55) we have that $(C_\lambda, 0)$ can be embedded in $(C^{g+1}, 0)$ by the sequence of approximate roots as in (13). By [G-T] the family $(C_\lambda, 0)$ admits simultaneous embedded resolution. This implies that $(C_\lambda, 0)$ is equisingular at $\lambda = 0$.

Suppose that the family $(C_\lambda, 0)$ is equisingular at $\lambda = 0$ to a plane branch $(C, 0)$. We have that F_λ is of the form (54). By Lemma 1.5 the approximate root $F_{j,\lambda}$ of F_λ of degree $n_1 \cdots n_j$ is of the form $F_{j,\lambda} = (y_1^{n_1} - \theta_1(\lambda) x_1^{m_1})^{n_2 \cdots n_j} + \cdots$, where the terms which are not written lie above the compact edge of the Newton polygon with respect to the coordinates (x_1, y_1). Then $F_{j,\lambda}$ defines an equimultiple family and that $y = 0$ is tangent to the curve $C_{j,\lambda}$ defined by $F_{j,\lambda} = 0$. By Proposition 1.8 we deduce that if $(C_\lambda, 0)$ is equisingular at $\lambda = 0$ then $C_{j,\lambda}$ is also equisingular at $\lambda = 0$, in particular the characteristic exponents with respect to $x = 0$ are constant in the family. The result is consequence of Theorem 4.1 for $F_{j+1,\lambda}$ with respect to $y = F_{1,\lambda}, F_{2,\lambda}, \dots, F_{j,\lambda}$. $\square$

We give an algorithm to check whether an equimultiple family of curves $(C_\lambda, 0)$ defined by a polynomial of the form (53) is equisingular at $\lambda = 0$ to a plane branch $(C_0, 0)$. We do not need to know the invariants of $(C_0, 0)$ a priori. If the answer of the algorithm is NO, then either $(C_0, 0)$ is not analytically irreducible or $(C_\lambda, 0)$ is not equisingular at $\lambda = 0$.

Algorithm 4.7. (1) *Step 1.*
    (a) *Set* $E_0 := N$, $\bar{B}_1 := (F_0, x)$ *and* $N_1 := \gcd(E_0, \bar{B}_1)$.
    (b) *Compute the approximate root* $F_{2,\lambda}$ *of degree* N_1 *of* F_λ.
    (c) *Check that* $F_{2,\lambda} = y^{N_1} - \theta_2(\lambda) x^{M_1} + \sum_{rN_1 + sM_1 > N_1 M_1,}^{s < N_1 - 1} \theta_{r,s}(\lambda) x^r y^s$, *where* $\gcd(N_1, M_1) = 1$ *and* $\theta_1(0) \neq 0$.
(2) *Step 2.*
    (a) *Compute* $\bar{B}_2 := (F_0, F_{2,0})_0$. *Set* $E_2 := \gcd(E_1, \bar{B}_2)$ *and* $N_2 := E_1 / E_2$.
    (b) *Compute* $F_{3,\lambda}$ *the approximate root of* F_l *of degree* $N_1 N_2$.

(c) *Compute the* (x, F_1, F_2)*-expansion of* $F_{3,\lambda}$.

(d) *Check that* $F_{3,\lambda} = F_{2,\lambda}^{N_2} - \theta_3(\lambda) M_2^{(3)}(0, M_2) + \sum_{rN_2+sM_2>N_2M_2}^{s<N_2-1} \theta_{r,s}(\lambda) M_2^{(3)}(r,s)$, *where* $\gcd(N_2, M_2) = 1$ *and* $\theta_2(0) \neq 0$.

(3) *Iterate Step 2.*

The algorithm stops whenever some condition is not verified answering NO or when $E_G = 1$ *for some integer* G *and then the answer is YES.*

To apply step 2 notice that by Remark 2.5 the exponents of a term $x^{i_0} F_{1,\lambda}^{i_1} F_{2,\lambda}^{i_2}$ appearing in the $(x, F_{1,\lambda}, F_{2,\lambda})$-expansion of $F_{3,\lambda}$ together with N_1, N_2, M_1 determine the integers (r, s) such that $x^{i_0} F_{1,\lambda}^{i_1} F_{2,\lambda}^{i_2} = M_2^{(3)}(r, s)$.

4.3.1. *Equisingularity criterion by jacobian Newton polygons.* Let $f \in \mathbf{C}\{x\}[y]$ be a Weierstrass polynomial. The *jacobian Newton polygon* of f with respect to the line $x = 0$ is the Newton polygon of $\mathcal{J}_f(t, x) := \mathrm{Res}_y(t - f, \frac{\partial f}{\partial y})$, where Res_y denotes the resultant with respect to y. The jacobian Newton polygon appears in more general contexts related to invariants of equisingularity (see [T1] and [T2]). García Barroso and Gwoździewicz have proved that if $f' \in \mathbf{C}\{x\}[y]$ is irreducible and $\mathcal{J}_f(t, x) = \mathcal{J}_{f'}(t, x)$ then f is irreducible. They have given two methods which characterize jacobian polygons of plane branches among other Newton polygons by a finite number of combinatorial operations on the polygons (see [GB-G], Theorem 1, 2 and 3). The following algorithm is consequence of their work.

Algorithm 4.8. *Imput: A family* $F_\lambda(x, y)$ *of the form (53).*

1. *Compute* $\mathcal{J}_{F_\lambda}(t, x)$.
2. *Compute the Newton polygon* $\mathcal{N}_\lambda$ *of* $\mathcal{J}_{F_\lambda}(t, x)$.
3. *Check that* $\mathcal{N}_\lambda = \mathcal{N}_0$.
4. *Test if* $\mathcal{N}_0$ *is a jacobian Newton polygon of a plane branch by using Theorem 2 or Theorem 3 in* [GB-G].

If all the steps of the algorithm give a positive answer then $F_\lambda = 0$ *is equisingular at* $\lambda = 0$ *to a plane branch.*

Acknowledgement. The author is grateful to J.-J. Risler for his remarks and suggestions on previous versions on this paper and to B. Teissier and P. Popescu-Pampu for useful discussions on this topic.

References

[A'C-Ok] A'Campo, N., Oka, M., Geometry of plane curves via Tschirnhausen resolution tower, *Osaka J. Math.*, **33**, (1996), 1003-1033.

[Abh1] Abhyankar, S.S. Inversion and invariance of characteristic pairs. *Amer. J. Math.* **89** (1967) 363–372.

[Abh2] Abhyankar, S.S., *Lectures on expansion techniques in algebraic geometry. Notes by Balwant Singh.* Tata Institute of Fundamental Research Lectures on Mathematics and Physics, 57. Tata Institute of Fundamental Research, Bombay, 1977.

[Abh3] Abhyankar, S.S., On the semigroup of a meromorphic curve. I. *Proceedings of the International Symposium on Algebraic Geometry (Kyoto Univ., Kyoto, 1977)*, pp. 249–414, Kinokuniya Book Store, Tokyo, 1978.

[Abh4] Abhyankar, S.S., Irreducibility criterion for germs of analytic functions of two complex variables. *Adv. Math.* **74** (1989), no. 2, 190–257.

[A-M] Abhyankar, S.S., Moh, T., Newton-Puiseux Expansion and Generalized Tschirnhausen Transformation I-II, *J.Reine Angew. Math.*, **260**. (1973), 47-83; **261**. (1973), 29-54.

[A-M2] Abhyankar, S.S., Moh, T., Embeddings of the Line in the Plane. *J.Reine Angew. Math.*, **276** (1975), 148-166.

[As-B] Assi, A., Barile M., Effective construction of irreducible curve singularities. *Int. J. Math. Comput. Sci.* **1** (2006), no. 1, 125–149.

[C] Campillo, A., *Algebroid Curves in positive characteristic*, Lecture Notes in Mathematics (813), Springer, Berlin, 1980.

[CN] Cassou-Nogués, Pi. Courbes de semi-groupe donné. *Revista Matemática de la Universidad Complutense de Madrid* **4**, (1991), no 1, 13–44.

[C-M1] Cossart, V., Moreno-Socías, G. Irreducibility criterion: a geometric point of view. *Valuation theory and its applications, Vol. II (Saskatoon, SK, 1999)*, 27–42, Fields Inst. Commun., 33, Amer. Math. Soc., Providence, RI, 2003.

[C-M2] Cossart, V., Moreno-Socías, G. Racines approchées, suites génératrices, suffisance des jets, *Annales de la Faculté des Sciences de Toulouse* Vol XIV, no 3, 2005, pp. 353-394.

[GB] García Barroso, E.R., Sur les courbes polaires d'une courbe plane réduite. *Proc. London Math. Soc.* **3** 81 (2000), no. 1, 1-28.

[GB-G] García Barroso, E.R., Gwoździewicz, Characterization of jacobian Newton polygons of plane branches and new criteria of irreductibility. ArXiv: 0805.4257[math.AG]

[G-T] Goldin, R., Teissier, B., Resolving singularities of plane analytic branches with one toric morphism, *Resolution of Singularities, A research textbook in tribute to Oscar Zariski.* Edited by H. Hauser, J. Lipman, F.Oort and A. Quiros. Progress in Mathematics No. 181, Birkhäuser-Verlag, 2000, 315-340.

[GP] González Pérez, P.D., Toric embedded resolutions of quasi-ordinary hypersurface singularities. *Ann. Inst. Fourier (Grenoble)* **53** (2003), no. 6, 1819–1881.

[GP-R] González Pérez, P.D.; Risler, J.-J., Multi-Harnack smoothings of real plane branches *preprint*, 2007.

[G-P] Gwoździewicz, J., Ploski, A., On the Approximate Roots of Polynomials. *Annales Polonici Mathematici*, **LX** 3, (1995), 199-210.

[G] Gwoździewicz, J., Kouchnirenko type formulas for local invariants of plane analytic curves, available at arXiv:0707.3404v1 [math.AG]

[K] Kouchnirenko, Polyèdres de Newton et nombres de Milnor. *Invent. Math.* , **32** (1976), no. 1, 1–31.

[L-Ok] Lê D.T. and Oka, M.; On resolution complexity of plane curves. *Kodai Math. J.* **18** (1995), no. 1, 1–36.

[LJ] Lejeune-Jalabert, M., Sur l'équivalence des singularités des courbes algebroïdes planes (coefficients de Newton), *Introduction à la theorie des singularités I (singularités et monodromie).* Edited by Lê Dũng Tráng. Hermann Paris, 1988, 49-154.

[M] Merle, M., Invariants polaires des courbes planes. *Invent. Math.* 41 (1977), no. 2, 103–111.

[Ok1] Oka, M., Geometry of plane curves via toroidal resolution, *Algebraic Geometry and Singularities*, Edited by A. Campillo López and L. Narváez Macarro. Progress in Mathematics No. 139, Birkhäuser, Basel, 1996.

[Ok2] Oka, M.*Non-degenerate complete intersection singularity*, Actualités Mathématiques, Hermann, Paris, 1997.

[PP] Popescu-Pampu, P., Approximate roots. *Valuation theory and its applications, Vol. II (Saskatoon, SK, 1999)*, Fields Inst. Commun., 33, Amer. Math. Soc., Providence, RI, 2003, 285-321.

[R] Risler, J-J., Sur l'idéal jacobien d'une courbe plane, *Bulletin de la S. M. F.* **99** (1971), 305–311.

[T1] Teissier, B., Teissier, B., Variétés polaires. I. Invariants polaires des singularités d'hypersurfaces. *Invent. Math.* **40** (1977), no. 3, 267–292.

[T2] Teissier, B., Polyèdre de Newton jacobien et équisingularité. *Seminar on Singularities (Paris, 1976/1977)*, pp. 193–221, Publ. Math. Univ. Paris VII, 7, Univ. Paris VII, Paris, 1980.

[T3] Teissier, B., The monomial curve and its deformations, Appendix in [Z2]

[T4] Teissier, B., Complex curve singularities: a biased introduction. *Singularities in geometry and topology*, 825–887, World Sci. Publ., Hackensack, NJ, 2007.

[W] Wall, C. T. C. *Singular points of plane curves.* London Mathematical Society Student Texts, 63. Cambridge University Press, Cambridge, 2004.

[Z1] Zariski, O., Studies in equisingularity III, Saturation of local rings and equisingularity on Algebraic Varieties, *Amer. J. Math.*, **90**, 1968 961–1023.

[Z2] Zariski, O., *Le problème des modules pour les branches planes*, Hermann, Paris, 1986.

Departamento de Algebra. Facultad de Ciencias Matemáticas. Universidad Complutense de Madrid. Plaza de las Ciencias 3. 28040. Madrid. Spain.

E-mail address: `pgonzalez@mat.ucm.es`